



*Government of Malawi Ministry of Health*

---

## **Integrated HIV Program Report July -September 2014**

- *Integrated HIV Program Supervision*
- *HIV Testing and Counselling / Early Infant Diagnosis*
- *Blood Safety*
- *Post Exposure Prophylaxis*
- *HIV Exposed Child Follow-Up*
- *Pre-ART*
- *Prevention of Mother to Child Transmission /  
Antiretroviral Therapy*
- *TB / HIV*
- *Sexually Transmitted Infections*
- *Supply of HIV Program Commodities*

<b>1</b>	<b>EXECUTIVE SUMMARY .....</b>	<b>1</b>
<b>2</b>	<b>INTEGRATED HIV PROGRAM OVERVIEW .....</b>	<b>2</b>
<b>3</b>	<b>SUPPORTIVE SITE SUPERVISION .....</b>	<b>3</b>
3.1	METHODS.....	3
3.2	SUPERVISION OUTCOMES.....	3
<b>4</b>	<b>INVENTORY OF SITES AND SERVICES .....</b>	<b>5</b>
<b>5</b>	<b>HIV TESTING AND COUNSELLING PROGRAM OUTPUTS .....</b>	<b>6</b>
5.1	HTC ACCESS TYPE .....	6
5.2	AGE AND SEX DISTRIBUTION AMONG HTC CLIENTS.....	6
5.3	FIRST TIME, REPEAT AND CONFIRMATORY TEST RESULTS.....	6
<b>6</b>	<b>DNA-PCR TESTING FOR EARLY INFANT DIAGNOSIS OF HIV (EID).....</b>	<b>8</b>
<b>7</b>	<b>BLOOD SAFETY .....</b>	<b>8</b>
<b>8</b>	<b>POST EXPOSURE PROPHYLAXIS (PEP) .....</b>	<b>9</b>
<b>9</b>	<b>PROVIDER-INITIATED FAMILY PLANNING (PIFP).....</b>	<b>9</b>
<b>10</b>	<b>COTRIMOXAZOLE PREVENTIVE THERAPY (CPT) .....</b>	<b>9</b>
<b>11</b>	<b>TB / HIV INTERVENTIONS .....</b>	<b>10</b>
11.1	INTENSIFIED CASE FINDING (ICF).....	10
11.2	ISONIAZID PREVENTIVE THERAPY (IPT).....	10
<b>12</b>	<b>HIV-RELATED DISEASES .....</b>	<b>11</b>
<b>13</b>	<b>HIV-EXPOSED CHILD FOLLOW-UP .....</b>	<b>11</b>
13.1	METHODS AND DEFINITION OF INDICATORS .....	11
13.2	HIV EXPOSED CHILD REGISTRATION DATA .....	12
13.3	BIRTH COHORT OUTCOMES .....	12
<b>14</b>	<b>PRE-ART .....</b>	<b>13</b>
14.1	PRE-ART REGISTRATION DATA .....	13
14.2	CUMULATIVE PRE-ART FOLLOW-UP OUTCOMES .....	13
<b>15</b>	<b>PMTCT / ART .....</b>	<b>13</b>
15.1	DATA SOURCES AND REPORTING METHODS.....	13
15.2	ARV COVERAGE AMONG PREGNANT / BREASTFEEDING WOMEN AND EXPOSED INFANTS .....	15
15.3	HIV SERVICES AT ANC.....	16
15.4	HIV SERVICES AT MATERNITY.....	16
<b>16</b>	<b>ART ACCESS AND FOLLOW-UP OUTCOMES.....</b>	<b>17</b>
16.1	NEW ART REGISTRATIONS DURING Q3 2014.....	17
16.2	CUMULATIVE ART REGISTRATIONS UP SEPTEMBER 2014 .....	19
16.3	ART OUTCOMES .....	19
16.4	ART COHORT SURVIVAL ANALYSIS.....	21
16.5	VIRAL LOAD (VL) MONITORING .....	24
<b>17</b>	<b>TB / HIV MANAGEMENT.....</b>	<b>24</b>
<b>18</b>	<b>STI TREATMENT.....</b>	<b>25</b>
18.1	ACCESS TO STI TREATMENT AND COVERAGE .....	25
18.2	CLIENT TYPE AND STI HISTORY .....	26
18.3	HIV STATUS.....	26
18.4	STI SYNDROMES.....	26
18.5	REFERRALS.....	26
<b>19</b>	<b>SUPPLY OF HIV PROGRAM COMMODITIES .....</b>	<b>27</b>
19.1	QUANTIFICATION AND PROCUREMENT PLANNING .....	27
19.2	QUARTERLY DISTRIBUTION OF HIV COMMODITIES.....	27
19.3	QUARTERLY LOGISTICS MONITORING AND SUPPLY CHAIN TRAIL.....	27
19.4	NATIONAL STOCK STATUS OF HIV COMMODITIES.....	28
<b>20</b>	<b>TRAINING AND MENTORING .....</b>	<b>30</b>
<b>21</b>	<b>PARTICIPANTS IN Q1 2014 SUPERVISION (SITE VISITS 6 – 24 OCTOBER 2014) .....</b>	<b>31</b>
<b>22</b>	<b>APPENDIX (FULL NATIONAL HIV PROGRAM DATA).....</b>	<b>2</b>

# 1 Executive Summary

This is the 11<sup>th</sup> quarterly HIV Program report after implementation of the 2011 Integrated Clinical HIV Guidelines in July 2011. A summary of the key achievements between **July and September 2014** is provided below:

- Scale-up of integrated HIV services had reached the following number of sites:
  - **724** static (579 within and 145 outside of health facilities) and 188 outreach HTC sites
  - **707** (static) ART sites
  - **619** PMTCT sites (Option B+, all included in ART sites above)
  - **654** Pre-ART sites
  - **638** sites with HIV-exposed child follow-up
- **550,425** persons were tested and counselled for HIV; **178,882 (32%)** accessed HTC for the first time; **371,543 (68%)** were repeat testers and **10,495 (3%)** of these received confirmatory testing (after having tested positive in the past). This is equivalent to **35%** confirmatory testing coverage among 29,893 patients initiating ART this quarter. **32,347 (6%)** clients received a positive result for the first time.
- **15,528 (97%)** of 16,034 blood units collected were screened for (at least) HIV, hepatitis B and syphilis.
- **135,958 (89%)** of 153,514 women at ANC had their HIV status ascertained; **10,296 (8%)** of these were HIV positive. **130,167 (95%)** of 137,343 women at maternity had their HIV status ascertained; **9,893 (8%)** of these were HIV positive.
- **29,893** patients started ART this quarter.
- **521,319** patients were alive and on ART by end of September 2014. This means that **52%** of the estimated 1 million HIV positive population was on ART. <sup>1</sup> Estimated ART coverage among people in need for treatment<sup>1</sup> was **41%** (45,124 / 111,000) for children (<15 years) and **69%** (476,195 / 687,000) for adults.
- **78%** of adults and **78%** of children were retained alive on ART at 12 months after initiation.
- **454,763 (93%)** of 488,020 patients on first line adult ART were on regimen 5A (TDF/3TC/EFV).
- **11,283 <sup>2</sup> (85%)** of an estimated **13,317 <sup>1</sup>** HIV infected pregnant women in Malawi were on ART this quarter. **6,220 (52%)** of these were already on ART when getting pregnant and **5,063 (48%)** started ART during pregnancy/delivery.
- An additional **1,655 <sup>2</sup>** breastfeeding women started ART due to **Option B+** (in WHO stage 1/2)
- **78%, 73%, 69%** and **67%** of women started under **Option B+** were retained on ART at **6, 12, 24** and **36 months** after initiation, respectively.
- **9,087 (7%)** of infants discharged alive from maternity were known to be HIV exposed, **8,482 (93%)** of these received ARV prophylaxis (nevirapine). **6,666 (73%)** were enrolled in exposed child follow-up before age 2 months.
- A total of **10,613** HIV exposed children and **6,941** pre-ART patients were enrolled for follow-up in *HIV Care Clinics (HCC)* during this quarter.

---

<sup>1</sup> 2014 Spectrum estimates based on 2014 definition of eligibility for ART in Malawi (CD4<500, Option B+, UT for U5).

<sup>2</sup> Adjusted for double counting due to patient transfers / 'failed ART initiations' among women lost to follow-up within 6 months of ART registration.

## 2 Integrated HIV Program Overview

Malawi implemented a revised HIV Program in all health facilities following the release of the **2011 Malawi Integrated Clinical HIV Guidelines**. The second edition of these guidelines was published in March 2014 and implementation of revised policies commenced in April 2014. Key program policies include:

- **PMTCT Option B+**: Universal life-long ART for all HIV infected pregnant and breastfeeding women regardless of clinical or immunological stage.
- Standard **HIV exposed child follow-up** to age 24 months. This program aims to improve early infant diagnosis and ART initiation using DNA-PCR testing for all infants from age 6 weeks; rapid antibody testing is considered diagnostic from age 12 months and repeated at 24 months. HIV exposed child enrolment and follow-up should be integrated with maternal ART follow-up (*Option B+*) to improve retention and adherence.
- **Early ART initiation**: universal ART for children under 5 years (confirmed HIV infection, CD4% no longer required), children over 5 years and adults with a **CD4** count  $\leq 500$ , patients with HIV and hepatitis B co-infection.
- Transition to a **new first line ART regimens** for adults (Malawi regimen 5A: tenofovir / lamivudine / efavirenz) and children (Malawi regimen 2: zidovudine / lamivudine / nevirapine), including provision of paediatric ARV formulations for all children under 25kg. Transition to 5A was completed by end 2013.
- Standardized **pre-ART services** for all HIV-infected persons not yet eligible for ART. The pre-ART program aims to reduce the incidence of HIV-related diseases and to enable early ART initiation based on the new CD4 cell threshold (500) through scheduled CD4 count monitoring.
- Provider-initiated provision of **contraceptives and condoms** for all adults in pre-ART and ART clinics to reduce the rate of unwanted pregnancies among HIV-infected adults and to reduce HIV-transmission between sexual partners.
- Isoniazid preventive therapy (**IPT**) for pre-ART patients to reduce the incidence of TB and intensified TB case finding (**ICF**) for all patients in pre-ART and ART follow-up to enable early diagnosis and treatment of TB and to reduce TB transmission in HIV clinics.
- Roll-out of scheduled **viral load monitoring** to improve early detection of treatment failure and initiation of second line ART.

Implementation of PMTCT Option B+ requires provision of ART services at all health facilities with Maternal and Child Health services. This required a massive acceleration of the **decentralization of ART services** from 303 (static) sites by June 2011 to currently 707 sites.

## 3 Supportive Site Supervision

### 3.1 Methods

The Department for HIV and AIDS has coordinated quarterly supportive supervision visits to all health facilities with ART services since the start of the national treatment program in 2004. Supervision teams are composed of: experienced HIV clinicians; nurses and M&E staff from health facilities in the public and private sector; district and zonal PMTCT and ART coordinators; program officers and technical staff from the Department for HIV and AIDS; technical staff from implementing partners. The TB and HIV programs are working towards a full integration of their respective site supervision exercises.

Each quarter, a one day pre-supervision meeting is organised for all supervisors participating in the upcoming round to share program updates, discuss observations from the previous round, distribute materials and organise logistics, transport and accommodation.

Standard supervision forms are used to guide implementation of the supervision protocol, to update site information and collect M&E reports. Custom forms with previous data for each site are printed from the HIV Department database. The supervision forms include:

- Contact details of HIV service providers at each site
- Quality of service checklist
- Follow up on action points noted during the previous visit
- Next visit date
- M&E reports from HTC, ANC, maternity, exposed child and pre-ART follow-up, ART and TB
- Physical drug stock-level assessment
- Identification of sites in urgent need of clinical mentoring
- Semi-structured feedback and performance rating for the supervision teams by facility staff

One copy of the supervision form is returned to the Department for HIV and AIDS, where data are entered in a custom MS Access database to produce national reports and to manage program logistics and the commodity supply chain. A second copy of the supervision form is left at the sites.

The supervision protocol includes a systematic review and verification of primary records (patient cards and registers) at all sites. This effectively provides a quarterly quality audit for M&E records, which has resulted in exceptional accuracy and completeness of HIV Program data in Malawi. At the same time, the systematic chart review helps to identify complex cases or deviations from clinical protocol, allowing the supervision team to provide targeted mentoring and clinical advice. The quarterly supervision exercise also aims to boost staff morale and motivation through *Certificates of Excellence* that are awarded by MOH to sites with an excellent score on the quality of service checklist. A growing number of health workers from sites all over the country participate as supervisors in this quarterly exercise and this has strengthened the national HIV Program identity and has greatly facilitated communication between program staff at the national, zonal, district and facility level.

The HTC Program usually conducts a separate supportive site supervision exercise each quarter, targeting a sample of HTC sites both within and outside of health facilities. Supervision teams consist of district, zonal and national level HTC coordinators, supported by implementing partners.

### 3.2 Supervision Outcomes

**720** public and private sector facilities were visited for **clinical HIV program supervision** between 6<sup>th</sup> and 25<sup>th</sup> October 2014. The large number of sites was covered by **95** supervisors working in **23** teams that spent a total of **1,914 working hours** at the sites. Each site visit lasted on average **2.7** hours, but

up to 2 days were spent at the busiest sites. **318** sites were awarded a *Certificate of Excellence* for **excellent performance**. The number of sites with excellent performance increased from the previous quarter despite a more rigorous application of performance criteria. **79** sites had significant weaknesses and were rated to require **intensive mentoring**. The capacity to provide site mentoring will need to be further expanded.

**Table 1:** Outcomes of integrated HIV services supervision for 2014 Q3

Zone	Total facil. visited*	Supervision hours spent at facilities		Performance (# and % of sites)	
		Total	Average per site	Excellent perform.	Mentoring needed
NZ	127	337	2.7	51 40%	13 10%
CEZ	98	225	2.3	57 58%	11 11%
CWZ	160	455	2.9	58 36%	27 17%
SEZ	166	499	3	72 43%	15 9%
SWZ	169	398	2.4	80 47%	13 8%
<b>Malawi</b>	<b>720</b>	<b>1,914</b>	<b>2.7</b>	<b>318 44%</b>	<b>79 11%</b>

\* includes facilities that were visited for assessment of readiness, but that may have not (yet) been designated to provide integrated HIV services.

**Table 1** provides a summary of the supervision outcomes by zone. Most facilities were using the standard national M&E tools. **120** sites had cumulatively registered more than 2,000 ART patient and **42** of these had registered more than 5,000. **43 (36%)** of these high burden sites were using electronic data system for ART (EDS). Some NGO supported sites were using custom tools compatible with the national standard reporting requirements.

## 4 Inventory of Sites and Services

A total of **724** static sites reported HTC service provision in Q3 2014 and **145** of these were outside of health facilities. **188** outreach HTC sites.

**Table 2:** Facilities with integrated HIV services in the 5 Zones. Availability of services defined by performance (at least 1 patient enrolled) during 2014 Q3

Zone	Total fac.(1)	Facilities providing HIV services				CD4 count machines (2)			Results
		Exp. child	Pre-ART	PMTCT B+	ART	Installed	Functional		
NZ	130	115 88%	117 90%	107 82%	122 94%	32 25%	31 97%	3,035	
CEZ	97	90 93%	89 92%	87 90%	94 97%	17 18%	15 88%	2,333	
CWZ	163	132 81%	134 82%	131 80%	158 97%	32 20%	25 78%	4,362	
SWZ	164	143 87%	156 95%	140 85%	161 98%	44 27%	41 93%	13,639	
SEZ	163	158 97%	158 97%	154 94%	162 99%	58 36%	53 91%	8,189	
<b>Malawi</b>	<b>717</b>	<b>638 89%</b>	<b>654 91%</b>	<b>619 86%</b>	<b>707 99%</b>	<b>183 26%</b>	<b>165 90%</b>	<b>31,558</b>	

(1) Total facilities in the public / private sector designated to provide integrated HIV services in this quarter. Individual site selection is reviewed and may change each quarter.

(2) CD4 machines that have produced at least 1 result during the reporting period are defined as functional.

**Table 2** shows the distribution of the **717** sites designated to provide clinical HIV services in Q3 2014, by zone. At the national level, there were **707** (static) sites with at least one patient on ART, **619** sites had enrolled women under PMTCT Option B+; **654** sites were providing pre-ART services. The number of sites with pre-ART patients slightly increased from 647 in the previous quarter; largely due to new sites starting to provide services. **638** had enrolled HIV exposed children for follow-up. ART services were now available at almost all designated sites in the 5 zones. The SE had reached 99% of designated sites with ART services and 94% of designated sites with Option B+.

CD4 count machines (including 'point of care' machines) were installed at **183** sites, and **165 (91%)** of these had produced at least 1 result during Q3 2014. **31,558** CD4 results were produced in this quarter. 43% of these outputs were generated with 37 machines in the SW zone, implying that many CD4 machines continued to experience down-time or to be running considerably below capacity.

## 5 HIV Testing and Counselling Program Outputs

HTC protocols were revised in 2013 and a new HTC register was implemented in the course of a national re-training campaign for all HTC providers between May and November 2013. Protocol revisions include:

- Clear recommendations for re-testing based on the client's test result and risk assessment
- Proper documentation of confirmatory testing for clients with a prior positive result (usually performed at enrolment into care).

This is the second HTC report based on the 2013 HTC register. The full national HTC data are presented in the **Appendix**.

**550,425** people<sup>3</sup> were tested and counselled for HIV between July and September 2014. **525,248 (95%)** of these tests were performed at health facilities and **25,177 (5%)** were done outside of health facilities. **22,166 (4%)** of the facility based test were in facilities providing HTC only. Out of a total of **32,347** people newly diagnosed with HIV this quarter, **29,949 (93%)** were tested at health facilities, **1,869 (6%)** at stand-alone HTC sites and **529 (2%)** in community-based testing.

### 5.1 HTC access type

**275,014 (50%)** of people tested were patients receiving provider-initiated testing and counselling (PITC); **272,264 (49%)** accessed voluntary counselling and testing, door-to-door, community-based testing, etc.; **3,144 (<1%)** came for testing with a *Family HTC Referral Slip* (FRS) that was issued to a family member at a prior HTC encounter. Based on a total of **14,520** FRS issued to index clients this quarter, the successful referral rate for family members was **22%** (3,144/ 14,520). This is higher than previous quarter (16%). Referral slip issuance and utilization has remained low.

### 5.2 Age and sex distribution among HTC clients

Out of **550,425** people tested and counselled, **36%** were males and **64%** were females. **47%** of females were pregnant. The proportion of males (51%) and non-pregnant females (49%) was very similar, implying gender balanced access to HTC services. Pregnant women have to be excluded from the comparison of male and female access to HTC because testing among pregnant women is almost entirely provider-initiated and there is no comparable access route targeting males.

**53%** of all people tested and counselled were 25 years and above, **38%** were between 15-24 years and **9%** were children below 15 years. **121,609 (22%)** accessed HTC with their partners (as a couple).

### 5.3 First time, repeat and confirmatory test results

The 2011 and 2014 Malawi Clinical HIV Guidelines stipulate: *All patients need a confirmatory HIV test to rule out any possibility of mix-up of test results or fraudulent access to ART: either at enrolment into pre-ART follow-up, or before starting ART if the test to confirm was not done in pre-ART. Children under 12 months starting ART with a positive DNA-PCR do not need another confirmatory test before starting ART, but all need a confirmatory rapid antibody test at age 12 and 24 months.* This is the second quarter reporting on confirmatory test results as a proportion of those who are classified as repeat testers.

---

<sup>3</sup> Reports from the HTC register are based on client encounters. It is not possible to de-duplicate people who access HTC multiple times in the reporting period. However, very few individuals come for repeat testing in less than 3 months and the number of HTC encounters in one quarter is therefore assumed to represent individuals.

**178,882 (32%)** accessed HTC for the first time and **371,543 (68%)** were repeat testers. Based on the cumulative number of people who accessed HTC for the first time, a total of **5,426,018** people have been tested since introduction of the *first time HTC access* indicator in July 2007.


**32,347 (6%)** out of all clients received a positive result for the first time. Positive rapid test results among infants (**1,062**) and inconclusive test results (**1,795**) both accounted for **<1%** of new results given to clients.

**358,447 (96%)** of 371,543 repeat testers reported a *last negative* result. **10,495 (3%)** were reported as *previous positives* and all of these should have been classified as receiving a confirmatory test. For most of the **10,495 previous positives**, testing was probably initiated by a health worker before enrolment into care. However, *confirmatory test results* accounted for only **9,000 (86%)** of *previous positive* clients. The remainder (1,495) may have been misclassified as *new positive* or *new inconclusive* because they were among clients who independently sought confirmation of their positive status. **8,366 (93%)** of 10,495 confirmatory tests were concordant positive and **634 (7%)** were classified as *confirmatory inconclusive*. This category includes parallel concordant negative and discordant test outcomes (Determine HIV1/2 and Uni-Gold HIV1/2 are used in parallel for confirmatory testing). This relatively high proportion of clients who did not have a concordant positive confirmation may be explained by selective confirmatory testing among clients with doubts about their previous positive status, but it underscores the importance of both routine confirmatory testing before ART initiation as well as the need to continue strengthening HTC quality assurance processes.

The 10,495 confirmatory test results documented this quarter indicate that only **35%** of the 29,893 patients initiating ART this quarter received confirmatory testing and **Figure 1** shows that confirmatory testing coverage was low in all 5 zones. Only **72 (10%)** of facilities throughout the country had performed confirmatory testing for  $\geq 75%$  of patients newly initiated on ART. Implementation of the confirmatory testing policy will be further reinforced over the next quarters.

**Figure 1: Confirmatory HIV testing coverage at ART sites in the 5 zones**

Num.: total confirmatory HIV tests documented in HTC registers. Denom.: total new patients initiating ART at the site



## 6 DNA-PCR testing for Early Infant Diagnosis of HIV (EID)

DNA-PCR testing is performed at 7 labs (Mzuzu Central Hospital, Mzimba District Hospital, Kamuzu Central Hospital, Queen Elizabeth Central Hospital, DREAM Blantyre, Zomba Central Hospital and Partners in Hope, Lilongwe). EID counsellors collect infant blood samples as dried blood spots on filter paper. Health facilities are requested to maintain a standard EID DNA-PCR logbook to document EID samples and to track results. The logbook includes the dates of collection, dispatch, receipt of result from the lab and communication of the result to the mother. For the 4<sup>th</sup> time this quarter, supervision teams were asked to collect basic data from these logbooks.

**528 (83%)** of 638 sites with HIV exposed children in follow-up had collected and recorded at least 1 DNA-PCR sample during Q3 2014. A total of **8,419** DNA-PCR samples were collected and recorded. By the time the logbooks were reviewed (between 2 and 4 weeks after the end of the quarter), results had been received at the sites for **4,151 (49%)** of these specimens and **2,209 (53%)** of these results had been communicated to the mother. The proportion of results received at the sites was **73%, 55%** and **23%** for samples collected in July, August and September, respectively. A total of **129 (3%)** results received at the sites were positive.

Detailed results from the **7 laboratories** could not be included in this report due to delays with the data analysis at the MOH, Department of HIV and AIDS.

## 7 Blood Safety

The Malawi Blood Transfusion Service (MBTS) is striving to provide blood products for the entire country using voluntary non-remunerated donors and quality assured screening for transfusion transmissible infections (TTIs). For the last years, MBTS has not been able to meet the national demand and several hospitals continue to supplement or rely entirely on blood units collected from replacement donors. Complete reports from MBTS have been available throughout, but blood safety reports from health facilities have not been consistently available and it has been challenging to compile national reports relying on the data passively submitted by the sites. Therefore, the PMTCT/ART supervision teams were tasked with active collection of blood donor and cross-matching data from all visited health facilities. Some of the visited laboratories were not using the standard MOH registers and the aggregation of data for reporting may have been affected by incomplete documentation at some sites.

A total of **16,034** blood units were collected in Malawi during Q3 2014. MBTS collected **10,387 (65%)** of these, **100%** of which were screened comprehensively for the relevant TTIs (HIV, Hepatitis B, Hepatitis C, syphilis, malaria). In addition, **54** hospitals in Malawi collected a total of **5,647** units from replacement donors. **5,141 (91%)** of these units were screened for at least the 3 key TTIs (HIV, HepB and syphilis) and **1,237 (32%)** of these were also screened for HepC and malaria. This means that a total of **15,528 (97%)** of all 16,034 units collected by MBTS and from replacement donors this quarter were screened at least for HIV, HepB and syphilis. Based on the blood donor registers at the sites that collected blood from replacement donors, 505 were screened with any other combination of tests for TTIs.

A total of **8,421** potential replacement donors were documented in the blood donor registers at the facilities and 5,647 (67%) of these ended up donating. Facilities may have used different screening algorithms and potential donors may have been excluded on the basis of different criteria, including TTIs, blood group, haemoglobin concentration and/or clinical conditions. Testing for less prevalent TTs may have only been carried out for donors who passed the screening for more common conditions. In total, 83% of potential donors were tested for HIV, 82% for HepB, 82% for syphilis, 51%

for malaria and 26% for HepC. Detailed data on individual test outcomes among all potential blood donors are presented in the Appendix.

## 8 Post Exposure Prophylaxis (PEP)

A total of **1,106** persons received PEP during Q3 2014. This is a slight increase from the previous quarter (898).

## 9 Provider-Initiated Family Planning (PIFP)

The 2011 Integrated Clinical HIV Guidelines encourage health workers to routinely provide condoms to all adults in pre-ART and ART clinics. Women should also be offered at least the standard injectable contraceptive (Depo-Provera) during any pre-ART or ART visit. This policy aims to address the significant unmet need for family planning that had been observed among HIV patients in Malawi and to reduce the number of unwanted pregnancies among HIV-infected women (**PMTCT Prong 2**). HIV program reporting on PIFP is limited to women who received an injection of Depo-Provera in pre-ART and ART clinics during the last quarter. The report does not account for family planning need nor does it include women who accessed family planning services outside of HIV clinics.

**Table 3:** Number and % of women retained in HIV care \* who were on injectable contraceptives (Depo) by the end of 2014 Q3.

Zone	Pre-ART		ART		Both patient groups	
	Tot. women	On Depo	Tot. women	On Depo	Tot. women	On Depo
NZ	845	229 27%	29,561	6,684 23%	30,407	6,913 23%
CEZ	533	101 19%	24,103	2,895 12%	24,636	2,996 12%
CWZ	2,872	464 16%	60,917	10,110 17%	63,788	10,574 17%
SEZ	3,267	1,621 50%	92,924	42,376 46%	96,191	43,997 46%
SWZ	4,051	957 24%	100,082	23,086 23%	104,133	24,042 23%
<b>Malawi</b>	<b>11,568</b>	<b>3,371 29%</b>	<b>307,587</b>	<b>85,150 28%</b>	<b>319,155</b>	<b>88,521 28%</b>

\* estimated from the total number of patients retained in pre-ART and ART, multiplied by the proportions of females and adults registered

**Table 3** shows that **88,521 (28%)** of 319,027 women in care received Depo-Provera from HIV clinics in Q3 2014. The SW Zone had achieved the highest coverage among women in pre-ART and ART. PIFP access continued to be affected by stock-outs of Depo-Provera, but patient coverage and stock availability had improved this quarter with 537

(77%) of ART/PMTCT sites having stocks of Depo-Provera in October 2014.<sup>4</sup> This was mainly due to inclusion of Depo-Provera in the quarterly distribution of ARVs and other HIV commodities.

## 10 Cotrimoxazole Preventive Therapy (CPT)

All patients in HIV care are universally eligible for CPT in order to reduce the frequency and severity of several HIV-related diseases. Patients with confirmed HIV infection are provided lifelong CPT in pre-ART and ART clinics. CPT is also given to HIV exposed children until exposure to breast milk has stopped and HIV infection has been ruled out (usually around age 24 months). Fewer than 5% of patients are expected to require stopping of CPT due to toxicity.

**Table 4** shows that **606,151 (94%)** of 645,169 all patients in care were on CPT at the end of Q3 2014.

<sup>4</sup> Many Mission hospitals do not provide family planning.

**Table 4:** Number and % of patients retained in HIV care who were on cotrimoxazole and isoniazid preventive therapy (CPT, IPT) by the end of 2014 Q3.

Zone	CPT								IPT	
	Exp. child		Pre-ART		ART		All patient groups		Pre-ART	
	Tot. pat.	On CPT	Tot. pat.	On CPT	Tot. pat.	On CPT	Tot. pat.	On CPT	Tot. pat.	On IPT
NZ	7,558	5,706 75%	3,103	3,030 98%	52,322	51,572 99%	62,983	60,308 96%	3,103	2,537 82%
CEZ	7,342	5,768 79%	2,211	2,186 99%	41,925	41,692 99%	51,478	49,646 96%	2,211	1,720 78%
CWZ	15,388	13,413 87%	8,320	7,265 87%	105,424	103,153 98%	129,132	123,831 96%	8,320	6,079 73%
SEZ	29,398	25,650 87%	11,860	11,758 99%	148,965	146,020 98%	190,223	183,428 96%	11,860	10,949 92%
SWZ	27,431	23,352 85%	13,247	12,064 91%	170,675	153,523 90%	211,353	188,939 89%	13,247	10,015 76%
Malawi	87,117	73,889 85%	38,741	36,302 94%	519,311	495,959 96%	645,169	606,151 94%	38,741	31,300 81%

## 11 TB / HIV Interventions

### 11.1 Intensified Case Finding (ICF)

TB is one of the most important HIV-related diseases in Malawi and a considerable proportion of (mainly early) deaths on ART are attributed to undiagnosed TB. ICF is carried out using a standard symptom checklist at every HIV patient visit. ICF outcomes are documented on HIV exposed child, pre-ART and ART patient cards, but routine M&E reporting is currently limited to ART patients in order to reduce the burden of reporting secondary cohort outcomes. It is assumed that implementation of ICF is similar in pre-ART and exposed child follow-up.

**508,599 (98%)** of all patients retained on ART were screened for TB at their last visit before end of September 2014. As of that visit, **5,583 (1%)** patients were new TB suspects and had presumably been referred for examination by a clinician and for TB investigations. **1,667 (<1%)** patients had confirmed TB (clinical or lab based). Out of these, **1,528 (92%)** were confirmed to be on TB treatment and **139 (8%)** had not yet started or had interrupted TB treatment. An excerpt from the data in the **Annex (Cumulative ART outcomes)** is shown below.

#### Current TB status among ART patients (ICF)

ICF not done (Current TB status unknown/ not circ)	10,712	2%
ICF done	508,599	98%
TB not suspected	501,349	99%
TB suspected	5,583	1%
TB confirmed	1,667	0%
TB confirmed, not on treatment	139	8%
TB confirmed, on TB treatment	1,528	92%

### 11.2 Isoniazid Preventive Therapy (IPT)

All pre-ART patients with a negative screening outcome for TB symptoms are eligible for IPT. The first (large scale) distribution of isoniazid and pyridoxine for the HIV programs reached the sites during July 2012. **31,300 (81%)** of 38,741 patients retained in pre-ART were on IPT by the end of September 2014. Isoniazid was in stock at 649 facilities during the October 2014 supervision visit. IPT coverage may increase further over the next quarters.

## 12 HIV-Related Diseases

**Table 5** shows the number of patients treated for key HIV-related indicator diseases. **4,692** patients were started on TB treatment this quarter and HIV status was ascertained for **4,368 (93%)**. **2,149 (49%)** of these were HIV positive and **1,460 (68%)** of all HIV positives were already on ART when starting TB treatment. The Diflucan (fluconazole) donation program has received renewed attention and significant stocks of fluconazole were distributed to all sites in November 2012. In Q3 2014, **368** and **640** patients received Diflucan for acute cryptococcal meningitis and oesophageal candidiasis, respectively. **310** patients with Kaposi sarcoma were registered for ART in this quarter.

**Table 5:** Number new cases of key HIV-related diseases registered per quarter (KS = Kaposi Sarcoma, CM = cryptococcal meningitis, OC = oesophageal candidiasis).

	TB				KS *	CM *	OC *
	Tot. cases	HIV status asc.	HIV positive	Already on ART	Tot. cases	Tot. cases	Tot. cases
2013 Q4	4,526	4,110 91%	2,280 55%	1,538 67%	414	661	883
2014 Q1	4,342	3,903 90%	2,103 54%	1,431 68%	361	414	690
2014 Q2	4,258	3,773 89%	2,130 56%	1,619 76%	312	408	767
2014 Q3	4,692	4,368 93%	2,149 49%	1,460 68%	310	368	640

## 13 HIV-Exposed Child Follow-Up

### 13.1 Methods and Definition of Indicators

There are multiple entry points into HIV exposed child follow up: children of HIV infected mothers may be enrolled at birth at maternity / postnatal ward; they may be found at Under 1 or Under 5 Clinics through active screening for HIV exposure; they may be identified when presenting sick to OPD; or they may be seen with their mothers in ART follow-up. Although the targeted enrolment age is below 2 months, children may theoretically be enrolled up to 23 months of age (when HIV infection can be ruled out by rapid antibody test and breast milk exposure is likely to have stopped).

Initial registration data and details for every visit are recorded on an *Exposed Child Patient Card* and a subset of the registration data is copied in the *HIV Care Clinic (HCC) register* (one record per patient). Registration data are reported from the HCC register on a quarterly basis. Follow-up outcomes are reported monthly, selecting children who were **2, 12 and 24 months** old in the respective reporting month. Outcomes are determined from the latest visit details recorded on each card. HIV infection status is evaluated as **known negative** if a negative DNA-PCR or rapid test result was available at the last visit; HIV infection status is evaluated as **known positive** if a positive DNA-PCR result was available at any age or a positive rapid antibody test was available from age 12 months; HIV infection status is counted as **unknown** if HIV infection has not been confirmed and/or a negative test result pre-dated the last visit (assuming on-going HIV exposure through breast milk). All children under 24 months with confirmed HIV infection and those under 12 months with confirmed HIV infection through DNA-PCR or HIV antibody and symptoms of *presumed severe HIV disease* are **eligible for ART**.

The main outcome indicator for the HIV exposed child follow-up program is **HIV-free survival at 24 months of age**. This is defined as the proportion of children who were discharged as confirmed HIV uninfected by the age of 24 months.

## 13.2 HIV Exposed Child Registration Data

This is the 12<sup>th</sup> quarterly report from the standard follow-up program for HIV exposed children. **10,613** HIV exposed children were newly enrolled into follow-up during Q3 2014; **6,666 (63%)** of these were under the age of 2 months. This represents timely enrolment for **73%** of the 9,087 known HIV exposed children discharged from maternity this quarter. The total number of new enrolments (10,613) exceeds by 1,526 (17%) the total number of known HIV exposed children discharged from maternity (9,087). This apparent discrepancy may be explained by delayed enrolment of infants born in previous quarters; by double-counting of infants who transferred between sites; or by identification and enrolment of additional HIV exposed infants after birth. Overall, enrolment into follow-up for known HIV exposed infants appears to be almost complete.

The documentation of follow-up outcomes, particularly the updating of DNA-PCR results on patient cards, remained incomplete at several sites. This has led to an underreporting of ascertainment of HIV status among the 2 month old cohort.

## 13.3 Birth Cohort Outcomes

There were **7,236** infants in the **2 month age cohort**. **2,684 (37%)** had received a DNA-PCR result. **66 (2%)** of these were confirmed HIV infected. An additional **41** infants were diagnosed with *presumed severe HIV disease*, which means that a total of **107** infants were eligible for ART. **39 (36%)** of these had started ART. The proportion of positives starting ART has decreased from the previous quarter (53%). Out of the entire 2-month age cohort, **6,625 (93%)** were retained in exposed child follow-up, **39 (1%)** had started ART and **4 (<1%)** were discharged confirmed uninfected<sup>5</sup>. **16 (<1%)** were known to have died and **467 (7%)** had been lost to follow-up.

There were **8,184** children in the **12 month age cohort**. Current HIV infection status was known for **3,191 (39%)** children (DNA-PCR or rapid antibody test) and **168 (5%)** of these were confirmed HIV infected. **30 (<1%)** additional children had been diagnosed with *presumed severe HIV disease*, which means that a total of **198** children were eligible for ART. **161 (81%)** had started ART. Out of the entire age cohort, **5,550 (70%)** were retained in exposed child follow-up, **161 (2%)** had started ART and **41 (1%)** were discharged confirmed uninfected.<sup>5</sup> **2,110 (27%)** were lost to follow-up and **72 (1%)** were known to have died.

There were **7,268** children in the **24 month age cohort**. Current HIV infection status was known for **2,994 (41%)** children (DNA-PCR or rapid antibody test) and **183 (6%)** of these were confirmed HIV infected. **21** additional children had been diagnosed with *presumed severe HIV disease*, which means that a total of **204** children were eligible for ART. **181 (89%)** of these had started ART. Out of the entire age cohort, **878 (12%)** were retained in exposed child follow-up, **181 (3%)** had started ART and **2,641 (38%)** were discharged confirmed uninfected. **3,252 (46%)** were lost to follow-up and **88 (1%)** were known to have died.

**Confirmed HIV-free survival at age 24 months** in this quarter was only **38%**, which was implausibly low and related to the fact that only 41% in this cohort had a known HIV status. 4,274 (59%) children were classified as '*current HIV infection status unknown*' and many of these may be among the 3,252 children lost to follow-up and the 88 children who had died. However, 878 (12%) were retained in follow-up beyond age 24 months and a final rapid test was not available for these children, possibly due to continued breast feeding. There are still problems with scheduled HIV testing (and documentation of test results) at 6 weeks, 12 and 24 months of age.

---

<sup>5</sup> A small number of children may be rightfully discharged as 'confirmed uninfected' by 2 or 12 months of age, provided that HIV exposure through breast milk has definitely stopped (e.g. maternal death) and a negative HIV test was obtained at least 6 weeks thereafter.

## 14 Pre-ART

### 14.1 Pre-ART Registration Data

A total of **6,941** patients were newly registered for pre-ART follow-up in Q3 2014. **546 (8%)** of these were children aged 5-14 years. The number of new pre-ART enrolments continued to decline from the previous quarter (7,342 total, 596 children) due to the introduction of relaxed ART eligibility criteria in April 2014. Several sites had already established pre-ART services before July 2011 and the cumulative number of pre-ART patients ever registered was **176,725**.

### 14.2 Cumulative Pre-ART Follow-up Outcomes

**38,741 (23%)** of all patients ever registered were retained in pre-ART follow-up by the end of September 2014; **86,847 (51%)** had started ART; **43,127 (25%)** had been lost to follow-up; **1,859 (1%)** were known to have died. The proportion of patients starting ART is bound to increase in the cumulative pre-ART cohort analysis over time. Based on a subtraction of cumulative outcomes from the previous quarter, **4,533** pre-ART patients started ART during Q3 2014 and **4,512** were lost to follow-up and **41** died.

CPT coverage among pre-ART patients was **94%** in Q3 2014 while IPT coverage increased from 79% to **81%**. **3,220 (23%)** of 13,829 women had received Depo-Provera from their pre-ART clinic. Further details on CPT, Depo-Provera and IPT are presented in **Tables 3** and **4** in the sections above.

## 15 PMTCT / ART

The implementation of **PMTCT Option B+** has effectively integrated PMTCT and ART services. The program aims to initiate lifelong ART for all HIV infected women as early as possible in pregnancy. ART may be started and continued at ANC, labour and delivery, and at ART clinics. All infants born to HIV-infected women are supposed to start daily nevirapine prophylaxis for the first 6 weeks of life. Nevirapine syrup is given to women at ANC at the earliest opportunity to take home with instructions how to give it to the new-born.

### 15.1 Data Sources and Reporting Methods

New standard M&E tools for ANC and maternity were implemented in January 2010 and revised in Q2 2012 to reflect the Option B+ policy. ANC and maternity clinic registers and reporting forms include patient management information and all relevant data elements for the maternal and child health and HIV programs. The ANC register was specifically designed to avoid data duplication that previously affected PMTCT reports from ANC due to the inability to account for individual women's outcomes in the course of multiple visits. The cohort reporting system is designed to aggregate women's outcome data after they have completed their ANC visits. Data from ANC and maternity are collated and presented separately because records do not allow identification of individual women and hence are subject to double counting if not separated.

All patients starting ART are recorded using standard program monitoring tools (ART patient treatment cards and ART clinic registers). **ART baseline data** for all patients registered are reported each quarter from ART clinic registers. **ART outcomes** of all patients ever registered are reported after reviewing the cards of all new patients and of those who were on ART at the end of the previous quarter, updating the status of patients who have subsequently died, stopped or been lost to follow-up. Secondary outcomes such as current regimen, CPT status, side effects, adherence and TB status are reported for all patients retained on ART.

ART scale-up has resulted in a growing proportion of HIV-infected women who are already on ART when getting pregnant. Implementation of *Option B+* will further increase ART coverage in this

group. **Maternal ART coverage** is estimated from the number of pregnant women who were already on ART when getting pregnant (**maternity reports**) *plus* those who newly started ART when pregnant (**ART reports**).

**Maternity reports** capture ART status at the time of delivery (up to the time of discharge from the postnatal ward). The timing of ART initiation is categorized into: (any time) before pregnancy; during 1<sup>st</sup> / 2<sup>nd</sup> trimester; during 3<sup>rd</sup> trimester; during labour. About 97% of pregnant women in Malawi attend ANC, but only 83% of women in the general population deliver at a health facility in Malawi. Maternity reports therefore have the potential for undercounting the number of mothers and infants receiving ARVs. However, there is evidence from ANC and maternity reports that almost all of the known HIV infected women deliver at health facilities. ARV coverage among known positives is therefore reliably calculated from maternity reports. Women admitted at maternity who are referred to another facility before / after delivery are double-counted in aggregated maternity data. Assuming the probability of referral is independent of ART status, the number of women already on ART when getting pregnant is therefore **adjusted** by the overall proportion of referrals among women admitted to maternity.

**ART program reports** capture pregnancy (and breastfeeding) status at the time of *ART initiation*, providing information on the number of new women starting ART while pregnant (or while breastfeeding). ART reports do not capture women who become pregnant after starting ART. For the estimation of maternal ART coverage, the number of women starting ART in pregnancy is **adjusted for**:

**a) Double-counting** of women starting ART in pregnancy and subsequently transferring to another site. These women are counted multiple times as ‘pregnant at the time of starting ART’ in the quarterly ART cohort reports because the disaggregation of age, sex and reason for starting ART applies to all patients newly registered in the quarter, including transfers in. Separate *ART ‘survival’ analyses* are collected each quarter for women started under Option B+. The proportion of women transferred within 12 months of registration is used to adjust the quarterly number of pregnant women starting ART for transfers.

**b) Failed ART initiation** is thought to be the main underlying reason for early loss to follow-up among the Option B+ cohort. Patients are recorded on patient cards and in clinic registers when the first supply of ARVs is dispensed and all new entrants are counted as ART initiations in the quarterly ART cohort report. Recent operational studies indicate that most pregnant women lost to follow-up within the first 6 months never return after this first dispensing visit and many of these may have never actually started taking ART. The proportion of women lost to follow-up in the 6-month survival analysis is therefore used to adjust the number of pregnant women starting ART in the quarterly ART cohort reports for *failed initiations*.

**Infant PMTCT coverage** is estimated from maternity reports, based on the number of infants born to known HIV-infected women and discharged alive who started nevirapine prophylaxis.

Coverage is calculated by dividing the number of patients served by population denominators. The denominators are derived from expected pregnancies based on population projections and HIV prevalence from epidemiological surveillance (source: 2014 Spectrum model for Malawi). There are an estimated 12,425 HIV infected pregnant women in the population per quarter (1/4 of 49,700 in 2014).<sup>6</sup>

---

<sup>6</sup> 2014 Spectrum estimates based on 2014 definition of eligibility for ART in Malawi (CD4<500, Option B+, UT for U5).

## 15.2 ARV Coverage among Pregnant / Breastfeeding Women and Exposed Infants


**11,283 (85%)** of the estimated 13,317 HIV infected pregnant women in Malawi this quarter were on ART. This is based on **6,220**<sup>7</sup> women at maternity who were already on ART when getting pregnant and **5,063**<sup>8</sup> women who newly initiated ART in pregnancy.

An additional **1,655**<sup>9</sup> breastfeeding women started ART due to **Option B+** (in WHO clinical stage 1 or 2), bringing the total number newly started on ART under **Option B+** to **6,718**. Most women starting ART while breastfeeding were probably identified late in maternity or early in the postnatal period, but this group may also include some women who re-initiated after interrupting ART in pregnancy. **8,420** infants were confirmed to have started NVP prophylaxis at maternity.

**Figure 2** shows the transition from prophylactic ARV regimens for HIV infected mothers to universal ART under **Option B+** (registration data; not adjusted as above). The (less effective) single dose NVP regimen and AZT combination prophylaxis had been phased out by April 2012. The average number of pregnant women registered for ART each quarter **increased almost 6-fold** from **1,221** in the 12 month period before introduction of Option B+ to an average of **7,000** since Q4 2011.

**Figure 2: Transition from prophylactic ARV regimens for PMTCT to Option B+ in Malawi**

Women who moved to Option B+ from sdNVP / AZT were double counted between Q3 2011 - Q1 2012. It is likely that <12,000 total women were on ARVs during these quarters. Data on women already on ART when getting pregnant are only available from Q2 2012.



<sup>7</sup> 6,579 women who started ART before pregnancy admitted at maternity; reduced by 5.5% to adjust for double-counting of 7,486 referrals among 137,343 total admissions.

<sup>8</sup> 7,004 women registered at ART clinics who were pregnant at the time of starting ART; a) 8.6% are discounted to adjust for double-counting of transfers based on 712 of 8,318 women who transferred within 12 months of registration (12 month Option B+ survival analysis); b) 20.9% are discounted to account for presumed failed ART initiations based on 1,603 of 7,655 women lost to follow-up within 6 months of registration (6 month Option B+ survival analysis).

<sup>9</sup> 1,810 women registered at ART clinics who were breastfeeding at the time of starting ART; reduced by 8.6% to adjust for double-counting of transfers based on 712 out of 8,318 women who transferred within 12 months of registration (12 month Option B+ survival analysis). Failed ART initiations are thought to be less common among this group, so no further adjustment is made.

### 15.3 HIV Services at ANC

The full national data from ANC are presented in the **Appendix**.

**164,321** women attended ANC for their first visit between July and September 2014. This is 96% of the estimated 170,732 pregnant women in the 2014 population during one quarter.<sup>10</sup>

The following report covers the outcomes of the **153,514** women who started ANC between January and March 2014 and who had finished ANC between July and September 2014. **14,086 (9%)** of these started ANC in their first trimester. **11,691 (8%)** were tested for syphilis at ANC and **257 (2%)** were syphilis positive. The low testing rate probably explains the higher (2%) than expected proportion (<1%) of positives as the testing was likely selective of those suspected to be positive. Only **34,502 (22%)** of women in this cohort attended the minimum of 4 focussed ANC visits.

#### 15.3.1 HIV Ascertainment at ANC

**135,958 (89%)** of ANC attendees had their HIV status ascertained. This is higher than the previous quarter (83%). Out of all women whose HIV status was ascertained, **9,317 (7%)** presented with a valid documented previous HIV test result and **126,641 (93%)** received a new HIV test result at ANC. A total of **10,296 (7.6%)** women were found HIV positive. This is lower than the estimated 11% HIV prevalence among pregnant women in the 2010 ANC sentinel surveillance survey but consistent with the latest Spectrum projections (7.8% HIV prevalence among pregnant women in 2014).<sup>6</sup>

#### 15.3.2 ARV Coverage at ANC

**9,550 (93%)** of (known) HIV infected women attending ANC received ART. This represents **72%** coverage of the estimated 13,317 HIV positive pregnant women per quarter at the population level.

Of the **9,550** ANC women who were known to receive ART, **4,241 (44%)** were already on ART when starting ANC, **4,126 (43%)** initiated before 28 weeks of pregnancy and **1,183 (12%)** initiated during the last trimester of pregnancy. **9,517 (92%)** of HIV infected women at ANC were on Cotrimoxazole Preventive Therapy.

**8,036 (78%)** of known HIV infected women attending ANC received the infant dose of ARVs (nevirapine syrup) to take home.

### 15.4 HIV Services at Maternity

The full national data from maternity are presented in the **Appendix**.

Between July and September 2014, **129,857** women were admitted for delivery to maternity; **7,486** of these were referred to another facility before delivery, resulting in **137,343** total admissions to maternity during Q3 2014. Out of all admissions, **126,611 (96%)** delivered at health facilities, while **5,005 (4%)** had already delivered before reaching a facility. The **126,611** facility deliveries represent **74%** of the estimated 170,732 quarterly deliveries in the population in 2014 which is less than the 83% reported in the Integrated Household Survey Report of 2010-2011.

A total of **123,481 (96%)** deliveries were conducted by skilled birth attendants, **844 (1%)** by paramedical staff and **4,854 (4%)** were not attended by any of the above (probably mainly among women who delivered before reaching maternity). **16,489 (12%)** of women developed obstetric

---

<sup>10</sup> Estimated as ¼ of 682,926 births projected for 2014.

Malawi National Statistics Office. (2008). Malawi Population Projections 2008-2030.

Retrieved from [http://www.nsomalawi.mw/images/stories/data\\_on\\_line/demography/census\\_2008/MainReport/The\\_maticReports/Population\\_Projections\\_Malawi.pdf](http://www.nsomalawi.mw/images/stories/data_on_line/demography/census_2008/MainReport/The_maticReports/Population_Projections_Malawi.pdf)

complications. The most common leading complications were obstructed / prolonged labour (**5,592** cases) and post-partum haemorrhage (**2,775** cases). A total of **131,616** babies were born, **127,129 (97%)** were singletons and **4,487 (3%)** were twins/multiples. There were **129,284 (98%)** live births and **2,332 (2%)** stillbirths. **128,029 (99%)** of babies born alive were discharged alive and **1,255 (1%)** died before discharge. **129,068 (>99%)** of women were discharged alive and **111 (<1%)** women died before discharge, which is equivalent to a maternal mortality ratio of **85 per 100,000** live births among women attending maternity.

#### 15.4.1 HIV Ascertainment at Maternity

**130,167 (95%)** women had their HIV status ascertained at maternity. Out of these, **127,052 (98%)** presented with a valid previous HIV test result and **3,115 (2%)** received a new HIV test result. A total of **9,893 (8%)** women were HIV positive and **120,274 (92%)** were negative. The **130,167** women whose HIV status was ascertained at maternity represent **76%** of the expected 170,732 women delivering in the population.

HIV exposure status was ascertained for **122,661 (96%)** out of 128,029 babies born and discharged alive. **9,087 (7%)** of these were born to a known HIV positive mother.

#### 15.4.2 ARV Coverage at Maternity

A total of **9,635 (97%)** of known HIV infected women admitted to maternity received ART. Out of these, **6,579 (68%)** had started ART before pregnancy, **1,602 (17%)** initiated ART during the 1<sup>st</sup> or 2<sup>nd</sup> trimester, **1,190 (12%)** initiated during the 3<sup>rd</sup> trimester and **264 (3%)** initiated ART at maternity.

A total of **8,420 (93%)** of 9,087 infants who were known HIV exposed and discharged alive started daily NVP prophylaxis at maternity. This represents **63%** coverage of the estimated 13,317 HIV exposed infants born in the population in this quarter.

## 16 ART Access and Follow-Up Outcomes

The full national data from the ART Program are shown in the **Appendix**.

### 16.1 New ART Registrations during Q3 2014

By the end of September 2014, there were **707 static ART sites** in Malawi, managed by government, mission, NGOs and the private sector. Out of these, **95** were ART facilities in the private sector, charging a nominal MK500 per monthly prescription of drugs per patient.

Implementation of the Malawi Integrated Clinical HIV Guidelines started in July 2011, triggering a massive surge in new ART initiations (see **Figure 3**). **29,893** patients initiated ART in Q3 2014 and **6,754** patients were registered as a transfer in (already on treatment; 18% out of all 37,082 clinic registrations). These numbers are similar to previous quarter.


Among all new registrations **35%** were males and **65%** females. **7,004 (29%)** of females were pregnant and **6,974 (>99%)** of these were started under **Option B+** in WHO stage 1 or 2, independent of their CD4 count. An additional **1,810** women in WHO stage 1 or 2 were started because of breastfeeding, bringing the total number of women registered as started under **Option B+**<sup>11</sup> to **8,784**.

---

<sup>11</sup> Universal ART for all HIV infected pregnant and breastfeeding women in WHO stage 1 or 2, independent of CD4 count

**Figure 3: Patients newly initiated on ART and total ART clinic registrations per quarter**

Total ART clinic registrations include patients who transferred between sites. This results in double counting of patients at the national level. For 'patients newly initiated on ART' every patient is only counted once.



A total of **25,101 (68%)** of all patients registered started in WHO stage 1 or 2 and **15,503 (62%)** of these started due a low CD4 count. **9,988 (27%)** of patients registered started in WHO stage 3 and **1,626 (4%)** started in stage 4.

**2,880** children were registered at ART sites in Q3 2014. **686** of these were registered under the expanded policy of universal ART for children aged 12-59 months in WHO stage 1 or 2, independent of CD4 count. **113** children started ART with presumed severe HIV disease, which was lower than the previous quarter (126). **123** infants in WHO stage 1 or 2 who started due to confirmed HIV infection through DNA-PCR, which is slightly higher than the previous quarter (107). Early paediatric ART access has remained below expectations, but the relatively low number is consistent with reduced transmission rates due to Option B+: considering that 9,087 HIV exposed infants were identified at maternity and assuming a 2% transmission rate among the 97% of HIV positive mothers at maternity who received ART (and 20% transmission in the 3% who did not receive ART)<sup>12</sup>, only about 231 of these known HIV exposed infants may have been infected perinatally during Q3 2014. However, considering the projected 1,025 new infant HIV infections in the 2014 population per quarter<sup>6</sup>, early infant treatment coverage remains low at an estimated **12%** (123 / 1,025). The most significant bottleneck for early infant treatment remains the identification of HIV infected pregnant / breastfeeding women.

**1,130 (3%)** out of all ART clinic registrations were patients with TB: **719 (2%)** had a current and **411 (1%)** a recent history of TB. **310 (1%)** of patients registered had Kaposi's sarcoma.

<sup>12</sup> UNAIDS Reference Group on Estimates Modelling and Projections (2011). Working paper on mother-to-child-transmission rates for use in Spectrum. Geneva, UNAIDS.

## 16.2 Cumulative ART Registrations up September 2014


By the end of September 2014, there were a cumulative total of **929,253** clinic registrations, representing **745,133 (80%)** patients who newly initiated ART and **174,411 (19%)** patients who transferred between clinics. **9,709 (1%)** out of all clinic registrations were patients who re-initiated ART after treatment interruption. Out of all registrations, **36%** were males and **64%** were females, **91%** were adults and **9%** were children (<15 years). Private sector clinics accounted for **28,369 (3.1%)** of total patient registrations.

## 16.3 ART Outcomes

**521,319 patients were alive on ART** by the end of September 2014. This number includes **2,008** patients who were assumed to be 'in transit' as of the 30<sup>th</sup> September 2014, based on the difference between **176,419** patients *transferred out* and **174,411** patients *transferred in* at the facilities around the country. This difference is explained by patients registered as a *transfer out* in the last 2 months of the quarter who have not yet arrived at their new site.

Out of the **745,133** patients ever initiated on ART, **521,319 (69%)** were retained alive on ART, **71,525 (9%)** were known to have died, **158,580 (21%)** were lost to follow-up and **3,418 (<1%)** were known to have stopped ART. An estimated **476,195** adults and **45,124** children (<15 years) were alive on ART by the end of September 2014.

**Figure 4 Patients alive on ART at the end of each quarter, stratified by size of facility (number of patients alive on ART)**




**Figure 4** shows the increase of patients alive on ART by the end of each quarter. The number of patients alive on ART **increased by 16,169** in Q3 of 2014. **Figure 4** also illustrates the ongoing decentralization of Malawi's ART program. From Q3 2011, the greatest increase in ART patient numbers was seen at sites with fewer than 500 patients alive on ART. By the end of September 2014, **50%** of the national ART patient cohort was in care at sites with fewer than 2,000 patients.

**Figure 5: Quarterly rates of ART drop out (ART stop, defaulters and deaths)**

Numerator: new ART stops, new defaulters and new deaths in the respective quarter


Denominator: total patients retained alive at the end of the previous quarter plus new patients registered in the respective quarter)



**Figure 5** shows the considerable decrease of ART drop-out rates since the start of the national program. There were **2,006** new deaths, **9,275** new defaulters, and **464** new ART stops in Q3 2014. This translates into a quarterly death rate of **0.4%** and a defaulter rate of **1.7%** among the patients alive and on treatment in this quarter. These rates were similar to the previous quarter. Based on previous operational studies, about half of the patients in stage 3 and 4 who are classified as lost to follow-up are thought to have died. There is also an indication that 10-15% of pregnant women who were registered as ‘initiated on ART’ under Option B+ may have never actually started taking ARVs due to inadequate preparation at ANC. Importantly, the ascertainment of loss to follow-up requires updating of patient treatment cards after analysis of the most recent dispensing visit. Any lack in rigour in this process will lead to a misclassification of patients who have been lost to follow-up as ‘retained alive on ART’.

By end of September 2014, a cumulative **71,525 (9%)** patients were known to have died **158,580 (21%)** were lost to follow-up and **3,418 (<1%)** were known to have **stopped ART**.

**Figure 6:** Patients starting ART in WHO stage 4 and deaths in the first 3 months after ART initiation. (Shown as proportions among new patients registered each quarter)




**Figure 6** shows the considerable decline in **early mortality** since the start of the program. In Q2 of 2006 11% of new patients died within the first 3 months after ART initiation. There has been a remarkable decline in early mortality and the lowest point thus far has been reached in Q4 2013. The decrease in early mortality is probably mainly due to earlier ART initiation (patients in WHO stage 2 with a CD4 count below the threshold or in stage 3). It correlates well with the decline in the proportion of patients starting ART in WHO clinical stage 4 from 25% in 2005 Q2 to **4%** in Q3 2014. Slight fluctuations in the calculated early mortality rates are mainly due to inconsistent classification of month of death at the sites with electronic patient record systems. The 2014 guidelines are expected to further reduce early mortality, as more patients will be started in WHO stage 1 and 2 (universal ART for HIV infected pregnant and breastfeeding women and children under 5 years).

## 16.4 ART Cohort Survival Analysis

A 12, 24, 36, 48, 60, 72, 84, 96 and 108-month '**cohort outcome survival analysis**' was conducted for patients registered in Q3 of 2005, 2006, 2007, 2008, 2009, 2010, 2011, 2012 and 2013, respectively. A separate 12-month cohort outcome analysis was conducted for children who were under 15 years at the time of ART initiation and who registered for ART in Q3 2013. For the 10<sup>th</sup> time, a further subgroup analysis was done for women who started ART under **Option B+** during Q3 2012, Q3 2013 and Q1 2014. **78% of adults** and **78% of children** were retained alive on ART after 12 months on treatment. This is similar to the previous quarter, but remains below the WHO target of 85%. **Figure 7** shows the continuous improvement of long-term treatment outcomes over time. **59%** and **52%** of patients registered 5 and 7 years ago had been retained alive on ART.

**Figure 7:** Group cohort survival analysis: Proportion of patients retained alive on ART 12, 24, 36, 48, 60, 72, 84 and 96 months after ART initiation



**6-month group cohort survival** outcomes were known for **8,164 (94%)** of the 8,707 women registered as having started ART under *Option B+* in Q1 2014.<sup>13</sup> This number represents 509 (7%) women who transferred out and are therefore double counted and **7,655 (94%)** patients not transferred. **5,992 (78%)** of these were retained at 6 months after registration. **1,603 (96%)** of those not retained were lost to follow-up, **23 (1%)** were known to have stopped ART and **46 (3%)** were known to have died.

**12-month group cohort survival** outcomes were known for **8,318** women registered as having started ART under *Option B+* in Q3 2013.<sup>13</sup> This number represents **712 (9%)** women who transferred out and are therefore double counted and **7,606 (91%)** patients not transferred. **5,544 (73%)** of these were retained at 12 months after registration. **2,062 (96%)** of those not retained were lost to follow-up, **23 (1%)** were known to have stopped ART and **63 (3%)** were known to have died.

**24-month group cohort survival** outcomes were known for **9,852 (94%)** out of the 10,443 women registered as having started ART under *Option B+* in Q3 2012.<sup>13</sup> This number represents **1,073 (11%)** women who transferred out and are therefore double counted and **8,779 (89%)** patients not transferred. **6,059 (69%)** of these were retained at 24 months after registration. **2,541 (93%)** of those not retained were lost to follow-up, **61 (2%)** were known to have stopped ART and **119 (4%)** were known to have died.

**2,689 (26%)** of the women in the 24 month *Option B+* survival cohort had initiated ART in the breastfeeding period and **2,897 (28%)** started in the third trimester / in labour; considering the 24 month median breastfeeding period in Malawi (2010 MDHS), more than half of the women in this cohort can be assumed to have stopped breastfeeding. The **69% retention rate at 24 months** after ART initiation confirms for the second time that a high proportion of women started under *Option B+* remain on ART beyond the cessation of breastfeeding.

The majority of women classified as lost to follow-up are likely to have stopped/ interrupted ART, but others will have transferred to another facility without notifying the previous site and the actual

<sup>13</sup> Group cohort survival analyses were not available from some sites with electronic data systems. 'Reason for starting' may be reclassified for some patients, leading to minor inconsistencies in patients included in group cohort survival analyses.

proportion retained on ART may be higher than reported. The 6-month retention rate is the same as in the previous quarter. These are satisfactory results. Most of the women lost to follow-up failed to return after their first visit and many of these may have never actually started ART due to inadequate counselling and preparation in the initial phase of implementation. The program is examining the different modes of delivery closely to further improve uptake and retention on *Option B+*.

### 6 month survival *OptionB+*

#### Survival and retention in ART program

\*

##### ART cohort registration group outcomes

Total ART clinic registrations	8,164	100%
Transfers out (double counted)	509	6%
Total not transferred out (patients in cohort)	7,655	94%
Total alive on ART	5,992	78%
Total not retained	1,663	22%
Defaulted	1,603	96%
Stopped ART	14	1%
Died	46	3%

### 12 month survival *OptionB+*

#### Survival and retention in ART program

\*

##### ART cohort registration group outcomes

Total ART clinic registrations	8,318	100%
Transfers out (double counted)	712	9%
Total not transferred out (patients in cohort)	7,606	91%
Total alive on ART	5,544	73%
Total not retained	2,062	27%
Defaulted	1,976	96%
Stopped ART	23	1%
Died	63	3%

### 24 month survival *OptionB+*

#### Survival and retention in ART program

\*

##### ART cohort registration group outcomes

Total ART clinic registrations	9,852	100%
Transfers out (double counted)	1,073	11%
Total not transferred out (patients in cohort)	8,779	89%
Total alive on ART	6,058	69%
Total not retained	2,721	31%
Defaulted	2,541	93%
Stopped ART	61	2%
Died	119	4%

## 16.4.1 Secondary outcomes of patients retained on ART

Secondary outcomes are known for the **519,311** patients alive on ART who remained at their sites at end of the quarter. They are assumed to be similar for the 2,008 patients *in transit*.

## ART Regimens

**513,600 (99%)** of patients were on first line and **5,240 (1%)** were on second line regimens; **471 (<1%)** were on non-standard regimens. Non-standard regimens are not necessarily substandard regimens and include patients continuing an ART regimen that was started outside Malawi, patients in research programmes and patients in specialist care.

Among patients on first line regimens, **25,580 (5%)** were on paediatric formulations and **24,401 (95%)** of these were on the new standard first line for children (regimen 2P: AZT/3TC/NVP).

By the end of September 2014, **454,763 (93%)** of patients on adult first line were receiving regimen **5A** (tenofovir / lamivudine / efavirenz). **26,686 (5%)** were on regimen 2A (zidovudine / lamivudine / nevirapine), which was the main alternative regimen for patients with stavudine side-effects before transition to regimen 5A and **1,925 (<1%)** were on regimen 1A (stavudine / lamivudine / nevirapine).

## Adherence to ART

Pill counts and the number of missed doses were documented for **514,069 (99%)** out of all patients retained on ART and **469,725 (91%)** of these were classified as >95% adherent in Q3 2014. Manual estimation of adherence from pill counts is practically difficult and classification can be misleading. The ART program has switched to a direct evaluation of doses missed in 2010 to improve on accuracy of adherence assessment and plausible adherence levels are recorded with this method. However, there have also been persistent challenges with the analysis of adherence levels at sites with Electronic Data Systems (EDS) and adherence data from several of these sites could not be included in this report.

## ART Side Effects

**515,292 (99%)** patients on ART had information on drug side effects documented at their last clinic visit before end of September 2014. **8,155 (2%)** of these had side-effects. The prevalence of side effects seems to have stabilized at very low levels following the full transition to regimen 5A (tenofovir / lamivudine / efavirenz) that started in July 2013.

## 16.5 Viral Load (VL) Monitoring

The National Treatment Program has started rolling out routine VL monitoring for patients on ART to facilitate early detection of treatment failure and timely switching to second line ART. Routine VL monitoring is scheduled at 6 months after ART initiation, at 2 years and every 24 months thereafter. Additional targeted VL testing may be carried out for patients with clinically suspected treatment failure. During Q3 2014, **8** laboratories in the national program provided VL testing for patients enrolled at the 8 respective facilities and associated sites.

Detailed results from these laboratories could not be included in this report due to delays with the data analysis at the MOH, Department of HIV and AIDS.

## 17 TB / HIV Management

Approximately **84%** of HIV infected TB patients were receiving ART in Q3 2014. This estimate is based on the following triangulation of TB and ART program data:

TB Program Data: A total of **4,692** TB patients were registered during Q3 2014. Assuming an average HIV prevalence of 60% among TB patients, **2,815** TB patients were HIV positive and therefore in need of ART. Given that **1,460** TB patients registered were already on ART at the time of starting TB treatment,  $2,815 - 1,460 = 1,355$  TB patients needed to initiate ART.

ART Program Data: An estimated **911** patients<sup>14</sup> started ART with a current or recent episode of TB in Q3 2014. This is **67%** (911 of 1,355) of the TB patients who needed to start ART. This means that a total of 1,460 + 911 = **2,371 (84%)** of the estimated 2,815 HIV infected TB patients were receiving ART in Q3 2014.

## TB program report

\*

### TB clinic registrations

Total TB patients registered	4,692	100%
------------------------------	-------	------

### HIV status ascertainment

HIV status not ascertained	324	7%
HIV status ascertained	4,368	93%
HIV negative	2,219	51%
HIV positive	2,149	49%
Already on ART	1,460	68%
Not on ART when starting TB treatment	689	32%

## TB / ART program triangulation

\*

### HIV-burden among TB patients (estimated)

HIV negative (est. 40%)	1,877	40%
HIV positive (est. 60%) in need of ART	2,815	60%
Not on ART	444	16%
Total on ART (coverage)	2,371	84%
Already on ART (TB prog)	1,460	62%
Started ART within 24m of TB diagnosis (ART prog)	911	38%
ART initiations with current TB (ART prog)	580	64%
ART initiations after recent TB (ART prog)	331	36%

## 18 STI Treatment

STI reports were actively collected during the Integrated HIV Program Supervision exercise for the 6<sup>th</sup> time this quarter. This decision was taken due to the persistent challenges with 'passive' reporting based on data aggregated by the district STI coordinators. The supervision teams noted that 78 (12%) of 650 facilities with STI services did not use the STI register (or used it inconsistently), so the data presented in this report are thought to represent 88% of STI clients treated at visited facilities. However, the *2013-14 Service Provision Assessment*<sup>15</sup> captured a total of 928 facilities offering STI management in Malawi and the site-level reports included here may therefore only represent 62% of all STI services in Malawi. The supervision teams re-emphasized the importance of complete and accurate documentation at the sites and the data quality is expected to improve with resumption of regular site supervision for the STI program. The complete set of STI program data collected is included in the Appendix.

### 18.1 Access to STI treatment and coverage

Based on the data collected at the facilities, a total of **54,427** STI cases were treated in Q3 2014. Considering the 88% completeness of reporting among the visited sites, this number is estimated to represent a total of **61,849** STI cases treated. This is equivalent to **63% STI treatment coverage** of the expected 98,600 STI cases in the population.

<sup>14</sup> 15% of the 1,107 ART patients who were registered this quarter with a recent or current episode of TB at the time of ART initiation were assumed to be transfers and were subtracted to adjust for double-counting.

<sup>15</sup> Ministry of Health, & ICF International. (2015). Malawi Service Provision Assessment (SPA) 2013-14. Lilongwe, Malawi and Rockville, Maryland, USA. Retrieved from <http://dhsprogram.com/pubs/pdf/SPA20/SPA20.pdf>

Out of **54,427** documented clients treated, **22,180** (41%) were male and **32,247** (59%) were female. **4,162** (13%) of female STI clients were pregnant. **37,014** clients (68%) were 25 years and above, **13,050** (24%) were 20-24 years and **4,363** (8%) were under 20 years old.

## 18.2 Client Type and STI History

**47,547** (87%) of clients were symptomatic and **6,880** (13%) were asymptomatic (treated as partners). Among symptomatic clients, **43,010** (90%) of were index cases and **4,537** (10%) were partners. A total of **12,507** partner notification slips were issued, equivalent to an average of 0.29 slips per index case. Considering the 12,507 partner notification slips issued, **91%** (11,417) of those notified presented to the clinic. **40,330** (74%) of clients presented with their first lifetime episode of STI, **9,006** (16%) clients reported to have had an STI in over three months ago and **5,091** (9%) of clients reported having had an STI within the last three months. Re-occurrence of an STI after a recent episode may be due to re-infection or treatment failure.

## 18.3 HIV Status

HIV status was ascertained for **27,119** (50%) clients and **7,175** (26%) of these were HIV positive. **1,495** (21%) of positives were identified through a new test initiated at the STI clinic, while **5,680** (79%) presented with a documented previous positive HIV test result. **4,196** (74%) of clients with a previous positive HIV test result were on ART.

The rate of HIV status ascertainment at STI clinics remained low. This is likely due to poor implementation of provider initiated testing and counselling, combined with weak back-referral systems which may lead to incomplete documentation of new HIV test results at the STI clinics. It is worth noting that a substantial proportion of clients who are aware of their HIV infection present with a new episode of an STI. This may suggest poor translation of positive living strategies promoted during counselling, but could also be due to the increased risk of recurrence of HSV-2 and balanitis among HIV-infected clients.

## 18.4 STI Syndromes

The most common syndrome was abnormal vaginal discharge (AVD) with **17,143** (29%) cases, followed by urethral discharge (UD, **13,474** cases) and genital ulcers (GUD, **9,770** cases). Similar to previous reports, balanitis, bubo, warts and neonatal conjunctivitis each accounted for 1 – 2% of cases.

## 18.5 Referrals

Given the high risk of recent HIV infection among STI clients, all clients with unknown status and those with a new negative test result should be referred for (repeat) HIV testing and counselling. **15,199** (31%) of the 54,427 STI clients with unknown or new negative test result were referred for repeat HTC. **880** (59%) of 1,495 clients who were newly tested HIV positive were referred for ART eligibility assessment.

## 19 Supply of HIV Program Commodities

### 19.1 Quantification and procurement planning

The quarterly quantification and procurement plan for all HIV commodities was reviewed and updated for the June 2014 order which included ARVs, OI medicines and selected laboratory commodities worth USD 70 million.

During Q3 2014, the supply situation for 5A (tenofovir / lamivudine / efavirenz) improved considerably compared with the previous quarters. The number of patients on regimen 5A increased by **18,039** (4.2%) from the previous quarter. **454,763 (93%)** of 488,020 patients receiving first line adult formulation ART were on this preferred regimen by the end of September 2014.

Almost universal availability of co-trimoxazole 960mg at sites (**653**) enabled the program maintain CPT coverage for 94% of pre-ART and 96% ART patients in Q3 2014.

During Q3 2014, ARVs and medicines for opportunistic infections worth \$33.8 million were received by the Central Medical Store Trust warehouse dedicated for HIV Program commodities. This included Regimen 5A (83% of the value of adult ARVs) and medicines for opportunistic infections (17% of the value for all medicines received during the period).

To maintain adequate stocks in the pipeline and hence ensure uninterrupted supply for subsequent orders, the Department of HIV and AIDS has continued processing HIV commodity orders for ARVs, OI, RDTs and other related commodities through Partnership for Supply Chain Management (ARVs and RDTs) and IDA Foundation (laboratory commodities and medicines for opportunistic infections). The first order for the transition funding period was placed in July 2014 to maintain adequate stocks between January and June 2015.

### 19.2 Quarterly distribution of HIV commodities

The scheduled quarterly distribution of HIV commodities (Distribution Round 18) took place between June and July 2014. 59 HIV commodities (ARVs, OI, STI medicines and laboratory commodities) were distributed over 700 ART sites. Both Determine and Unigold HIV test kits were also distributed to individual health facilities to ensure adequate stocks and uninterrupted testing services at all sites.

During Q3 2014, the logistics team at the Department of HIV and AIDS also coordinated a total of over 3,000 individual commodity transactions between sites to avert stock outs or expiries. All such transactions are all managed using the HIV Department Supply Chain Hot Line, a toll free facility that was set up to facilitate communication between the health facilities and the central level. Health workers are able to communicate supply chain and other drug related issues that need to be resolved by the technical team at the department in a timely manner.

### 19.3 Quarterly logistics monitoring and supply chain trail

Following distribution round 18, a sample of 32 facilities in Thyolo, Phalombe, Machinga, Ntcheu, Karonga, Mzimba North, and Nkhatabay were visited to physically verify receipts and to support facilities in their commodity management. No deviations were noted from the verified delivery notes reviewed by the team and health facility staff during the supply chain trail visit. Some of the challenges noted include poor documentation of emergency dispensations, highlighting the need to introduce a standard emergency dispensing register to improve the ability to account for all HIV commodities. There was a marked improvement in the management of ARVs and medicines for opportunistic infections. Health workers continued to use RDT daily activity registers to document utilization of every HIV test supplied. The standard registers for redistributed commodities with

authorization codes from the HIV Department were generally well filled. This tool has greatly improved the accountability for all HIV commodities issued to sites.

#### **19.4 National Stock Status of HIV Commodities**

Physical stock counts for ARVs and other medicines for HIV-related diseases were performed at all sites during the supervision visits in October 2014. **Table 6** shows the total medicine stocks found at the sites and the estimated consumption periods. Site-level stocks of the key adult and paediatric regimens were estimated to last until February 2015, which is consistent with maintaining a 3 month working stock plus a 2 month buffer stock at all service delivery points.

Minimum supplies of TDF/3TC 300/300mg and AZT/3TC 300/150mg are maintained at all sites for post-exposure prophylaxis (PEP) and the total stocks at the sites therefore far exceeds the actual consumption from patients using this regimen in alternative ART regimens or as PEP.

High volume commodities include stavudine- and efavirenz-containing regimens at all levels mainly driven by the transition to tenofovir based regimens. The program continues to monitor the trend of patients on stavudine and efavirenz containing regimens to inform future procurements.

The actual number of patients on 5A (454,763) deviated by only 1,695 (**0.37%**) from the forecast for end September 2014 (456,458). The national ART program forecast and quantification was updated in September 2014 to inform procurement planning and budgeting for HIV commodities.

**Table 6:** Total stocks of HIV program commodities at all sites visited during the 2014 Q3 supportive site supervision. Stock positions are from the date of the visit (between 1-4 weeks after the end of the quarter). Warehouse stock positions are from 26/01/2015

Inventory unit	Item	Sites with any Stock	Total Physical Stock		Consumption/ Month	Months of Stock *	
			At Sites	In Warehouse		At Sites	Wareh.
<b>tins</b>	ABC / 3TC 60 / 30mg tins (60 tabs)	119	19,051	8,605	3,657	5.2	2.4
	ABC / 3TC 600 / 300mg tins (30 tabs)	17	411				
	ATV / r 300 / 100mg tins (30 tabs)	153	20,192	21,183	4,582	4.4	4.6
	AZT / 3TC / NVP 300 / 150 / 200mg tins (60 tabs)	613	125,481	398,026	26,686	4.7	14.9
	AZT / 3TC / NVP 60 / 30 / 50mg tins (60 tabs)	627	342,540	568,892	61,003	5.6	9.3
	AZT / 3TC 300 / 150mg tins (60 tabs)	592	18,368	12,173	1,751	10.5	7.0
	AZT / 3TC 60 / 30mg tins (60 tabs)	587	20,498	15,521	2,229	9.2	7.0
	d4T / 3TC / NVP 30 / 150 / 200mg tins (60 tabs)	38	13,047	88	1,925	6.8	0.0
	d4T / 3TC 30 / 150mg tins (60 tabs)	55	9,762	97	158	61.8	0.6
	d4T / 3TC 6 / 30mg tins (60 tabs)	212	3,061		318	9.6	
	EFV 200mg tins (90 tabs)	118	2,192	1,367	330	6.6	4.1
	EFV 600mg tins (30 tabs)	387	20,846	15,117	843	24.7	17.9
	LPV / r 100 / 25mg tins (60 tabs)	53	8,308	9,870	1,974	4.2	5.0
	LPV / r 200 / 50mg tins (120 tabs)	57	1,120	222	458	2.4	0.5
	NVP 200mg tins (60 tabs)	430	22,904	52,562	3,723	6.2	14.1
	NVP 50mg tins (60 tabs)	108	9,261	13,350			
	TDF / 3TC / EFV 300 / 300 / 600mg tins (30 tabs)	702	1,961,555	2,402,319	454,763	4.3	5.3
TDF / 3TC 300 / 300mg tins (30 tabs)	615	43,882	56,753	7,977	5.5	7.1	
<b>bottles</b>	Fluconazole (Diflucan) 50mg / 5ml bottles (35 ml)	502	2,789		72	39.0	
	Gentian violet 25g bottles (1 each)	488	7,350		1,122	6.6	
	NVP 10mg/ml bottles (25 ml)	472	78,111	154,751	15,863	4.9	9.8
<b>vials</b>	Benzathine Penicillin 1.44g vials (50 each)	657	194,289		36,118	5.4	
	Bleomycine 15,000IU vials (1 each)	23	3,384				
	Ceftriaxone 1g vials (50 each)	556	138,651		97,489	1.4	
	Depo-Provera 150mg/1ml vials (25 each)	531	472,073	1,287,050	308,358	1.5	4.2
	Gentamicin 80mg / 2ml vials (50 each)	617	871,101		91,741	9.5	
	Vincristine 1mg / 1ml vials (1 each)	53	18,403	50,458	3,720	4.9	13.6
<b>tabs</b>	Aciclovir 200mg blister packs (25 tabs)	655	5,812,244	3,114,250	587,662	9.9	5.3
	Amitriptyline 25mg tins (500 tabs)	278	796,274		244,650	3.3	
	Azithromycin 500mg blister packs (3 tabs)	447	215,668		9,699	22.2	
	Ciprofloxacin 500mg blister packs (100 tabs)	380	986,516		277,973	3.5	
	Clotrimazole 500mg boxes (1 each)	467	61,779		35,731	1.7	
	Codeine 30mg tins (100 tabs)	189	110,836		46,006	2.4	
	Cotrimoxazole 100 / 20mg blister packs (1000 tabs)	629	32,137,611	49,703,000	5,616,506	5.7	8.8
	Cotrimoxazole 400 / 80mg tins (1000 tabs)	666	61,009,940	21,774,000	15,411,333	4.0	1.4
	Cotrimoxazole 960mg blister packs (1000 tabs)	653	43,191,436	20,936,000	16,406,729	2.6	1.3
	Doxycycline 100mg tins (1000 tabs)	614	21,153,702	998,000	4,118,702	5.1	0.2
	Erythromycin 250mg tins (1000 tabs)	440	6,199,512	1,498,000	3,684,596	1.7	0.4
	Fluconazole (Diflucan) 200mg tins (28 tabs)	395	436,095	291,760	35,198	12.4	8.3
	Fluconazole (generic) 200mg tins (100 tabs)	42	117,314				
	Ibuprofen 200mg tins (100 tabs)	427	3,766,273		787,775	4.8	
	Isoniazid 100mg blister packs (100 tabs)	153	231,865		141,327	1.6	
	Isoniazid 300mg tins (1000 tabs)	649	11,581,175	14,663,000	1,034,385	11.2	14.2
	Metronidazole 200mg tins (1000 tabs)	550	9,342,363		4,474,247	2.1	
	Morphine 10mg blister packs (60 tabs)	177	449,980		200,754	2.2	
	Pyridoxine 25mg tins (100 tabs)	122	1,242,566		1,104,119	1.1	
Pyridoxine 50mg tins (1000 tabs)	520	6,455,220	12,119,000	1,104,119	5.8	11.0	
<b>sheets</b>	ART pat. card adult (yellow) bundles (100 sheets)	679	217,307	23,900	11,401	19.1	2.1
	ART pat. card paed. (blue) bundles (100 sheets)	633	102,445		960	106.7	
	Exposed child card (pink) bundles (50 sheets)	634	69,919	49,250	3,538	19.8	13.9
	Polythene sleeve bundles (100 sheets)	564	131,476	294,600			
	Pre-ART pat. card (green) bundles (100 sheets)	628	147,995		2,314	64.0	
<b>tests</b>	DBS kit (filter paper, lancet, etc.) bundles (20 eac)	383	20,801	440	4,245	4.9	0.1
	Determine HIV1/2 boxes (100 each)	651	759,878	590,000	172,532	4.4	3.4
	Determine syphilis boxes (100 each)	53	14,276	90,200	51,120	0.3	1.8
	Uni-Gold HIV1/2 boxes (20 each)	545	41,766	14,060	13,654	3.1	1.0
<b>pieces</b>	Condoms female boxes (1000 each)	421	901,324		172,515	5.2	
	Condoms male boxes (144 each)	595	11,852,233	11,557,152	4,787,490	2.5	2.4

\* 'Consumption per month' and 'Months of stock' for ARVs, CPT, INH and HIV test kits are based on the respective patient-regimen groups in the standard service reports. Estimates are based on the number of patients on the respective regimen at the end of the quarter evaluated and do not account for potential (positive or negative) growth. Facility stock positions for OI and STI drugs include HIV Program and other supply sources. Total national consumption and MoS estimates are used for these commodity groups. 'Months of stock' is calculated from the day of the physical stock count, which is on average 1 month after the end of the quarter.

## 20 Training and Mentoring

### **Viral Load orientation trainings**

446 health workers were orientated in routine & targeted viral load testing using the capillary DBS sample technique.

### **ART/PMTCT refresher trainings:**

2,649 clinicians and nurses were refreshed on the new 2014 National guideline. Cumulatively, 4380 health workers (providers) have been refreshed.

### **Mentor of Mentor and Zonal Clinical Mentoring**

The mentor of mentor (MOM) training was conducted in the South west zone as one way enhancing the Clinical mentoring activities within the Zone. 4 districts participated. One designated MOM for each district who provides support to the mentors within the district.

The training reviewed the clinical mentoring activities in district and oriented the mentors on the revised monitoring and evaluation tools for the clinical mentoring program.

## 21 Participants in Q1 2014 Supervision (Site visits 6 – 24 October 2014)

Florence Nkonja (Nurse, MACRO)	Stuart Chuka (CO, MBCA)	Roseby Malombe (Nurse, CHAM)
Zengani Chirwa (TA, MOH, Department of HIV and AIDS)	Michael Eliya (PMTCT Program Officer, MOH)	Alefa Fikira (CMT, MOH)
A Phiri (Clerk, MOH)	Angela Nkhoma (Nurse, MOH)	Jesse Lobeni (Nurse, MOH)
Gift Kakwesa (, MOH)	Stanley Ngoma (CO, MOH)	Gerald Zomba (Program Officer, Dept for HIV and AIDS)
James Mwambene (CO, Diginitas)	Rodrick Kaulele (CO, CHAM (Sister Tereza))	Kenneth Matumba (, CHAM)
Chimwemwe Francis Mkandawire (IT Fellow, Dept for HIV and AIDS)	Mathilda Kamanga (Nurse, Army)	Eviness Kafumbi Nkhoma (Nurse, MOH)
Tewodros Wubayehu (Zonal Supervisor, CW Zone)	Timothy Mwenyedini (MA, MOH)	Sidder Hambisa (ENM, MOH)
Henry Banda (CO, MOH)	Marion Chikuse (, moh)	Ekwala Mubiala (HIV Zonal Supervisor, MOH, UNV)
Lameck Manda (Logistics Fellow, MOH)	Andrew Mganga (M&E Fellow, Dept for HIV and AIDS)	Dereje Habte (ART Zonal Supervisor, CE zone)
Harrison Tembo (CO, MOH)	Overtone Ndhlovu (CO, MOH)	Peter Donda (CO, Dedza DH)
Miriam Chigwiya (CO, MOH)	Joel Kazembe (CO, MOH)	Loyna Mbewe (Nurse, MOH)
Melenia Nkhoma (Logistics Fellow, MOH)	Limbani Kadzuwa (Nurse, MOH)	Ormisher Joe Nthala (CO, Lighthouse)
Rose Banda (, MOH)	Austins Namondwe (CO, CHAM)	Erik Mittochi (CO (ART coord), MOH)
Linly Mulundira (, lighthouse)	Peter Chimphero (CO, MOH)	Mercy Makaika (Nurse, MOH)
Chikayiko Majamanda (Nurse, MOH)	Lincy Chalunda (CO, MOH)	Davie Maseko (CO, SOS)
Cecilia Sambakunsi (Logistics Fellow, HIV Dept)	Peter Mwamulima (, Partners in Hope)	Damison Msiska (CO, Dwangwa)
Weston Njamwaha (Clinician, PIH)	Judith Ntopa (Nurse, Army)	Cecilia Manyawa (Nurse, MOH)
Monica Simfukwe (Nurse, MOH, Chintheche RH)	Moreen Mtambo (PMTCT, MOH)	Andraida Mtoseni (Nurse, MOH)
Fainala Muyila (Nurse, MOH)	Oscar Kasiyamphanje (Nurse, CHAM)	Sabina Phiri (Nurse, MOH)
Grace Chipanga (Nurse, Private)	Ruockia Mwachumu (Nurse, MOH Nsanje DHO)	Martin Katanga (CO, MOH)
Mapayi Ngalala (HIV Zonal Supervisor, MOH, UNV)	Amos Makwaya (CO, MOH)	Erick Mtemang'ombe (CO, CHAM)
Everista Mkandawire (Nurse, MOH)	Juliana Soko (ARV nurse, MOH, Livingstonia MH)	Mike Nyirenda (CO, Lighthouse)
Vera Kajawo (Nurse, MOH)	Hannock Matupi (ARV clinician, MOH, Rumphu DH)	William Mkandawire (NMT, CHAM)
Rose Maviko (Nurse, Limbe HC)	Ezra Majoni (Nurse, MOH)	Savior Mwandira (, PIH Lilongwe)
Edith Taulo (Nurse, MOH)	Beatrice Malonje (Nurse, MOH)	Mervis Ngonga (Nurse, MOH)
Peter Kamanga (CO, EQUIP)	Janet Chikonda (Nurse, MOH)	Eustice Mhango (ART officer, MOH, Department of HIV and AIDS)
Yusuf Bhamu (HIV Fellow, HIV Dept)	Dalitsa Midiani (PMTCT Officer, MOH)	Felix Chinguwo (CO, Ntcheu DH)
Jean Kayamba (Nurse, MOH)	Afred Kamoto (Logistics Fellow, MOH)	Simon Makombe (ART officer, MOH, Department of HIV and AIDS)
John Kabichi (CO, MOH)	Ruth Deula (Nurse, CHAM)	Grace Juma Nkhata (Nurse, MOH)
Lilian Kachali (Nurse, MOH)	Charles F Sekani (CO, .)	Absalom Kaunda (CO, MOH, Mzimba DHO)
Macleod Piringu (ART CORDINATOR, MOH)	Chiukepo Longwe (CO, Private)	Salome Chiwewe (Nurse, MOH, Ntchisi DH)
Cecelia Tenesi (Nurse, MOH)	Chifundo Makuluni (Nurse, MOH)	
	Augustine Mphweya (, PIH North)	

### **Report compiled by:**

Frank Chimbwandira (MO/Head of Dept. for HIV and AIDS)  
 Austin Mnthambala (MO/Deputy of Dept. for HIV and AIDS)  
 Simon Makombe (ART Officer, Dept. for HIV and AIDS)  
 Eustice Mhango (ART Officer, Dept. for HIV and AIDS)  
 Michael Eliya (PMTCT Officer, Dept. for HIV and AIDS)  
 Dalitsa Midiani (PMTCT Officer, Dept. for HIV and AIDS)

Lucius Ng'omang'oma (HTC Officer, Dept. for HIV and AIDS)  
 Andreas Jahn (MO/TA, Dept. for HIV and AIDS)  
 Zengani Chirwa (MO/TA, MOH, Dept. of HIV and AIDS)  
 Caroline Ntale (TA Logistics, MOH, Dept of HIV and AIDS)  
 Andrew Mganga (M&E Fellow, Dept. for HIV and AIDS)  
 Gerald Zomba (Clin. HIV Fellow, Dept. for HIV and AIDS)  
 Chimwemwe Mkandawire (IT Fellow, Dept. for HIV and AIDS)

We thank all facility staff for their sincere welcome and co-operation with the HIV Department and its partners during these supportive visits. We congratulate all staff for their excellent work.

30 January 2015

## 22 Appendix (Full National HIV Program Data)

# HTC site report

Malawi (national)

2014 Q3 (1st month of quarter, 2nd month of quarter, 3rd month of quarter)

## Clients at health facility (static)

### HTC client details

\*

#### Total HTC clients served

Total HIV tested	503,132	100%
------------------	---------	------

#### Sex

Males tested	174,902	35%
Females tested	328,230	65%
Females non-pregnant	168,123	51%
Females pregnant	160,107	49%

#### Age

Children 0-14 yrs	45,715	9%
Children below 12 mths (Age group A)	1,576	3%
Children 18 mths - 14 yrs (Age group B)	44,139	97%
Adults 15+ years	457,417	91%
Young adults 15-24 years (Age group C)	193,134	42%
Older adults 25+ yrs (Age group D)	264,283	58%

#### HTC access type

PITC	262,412	52%
Family Referral Slip (FRS)	3,067	1%
Other (VCT, etc.) HTC access	237,653	47%

#### HTC first time / repeat

Never tested before	162,549	32%
Previously accessed HTC	340,583	68%
Last negative	328,180	96%
Last positive	10,044	3%
Last exposed infant	1,965	1%
Last inconclusive	394	0%

#### Counseling session type / Partner present

Counseled with partner / partner present	116,135	23%
Counseled alone / Partner not present	386,997	77%

#### Outcome summary (HIV test)

Single test negative	462,175	92%
Single test positive	697	0%
Test 1&2 negative	544	0%
Test 1&2 positive	38,216	8%
Test 1&2 discordant	1,500	0%

#### Final result given to client

Results among clients never tested / last negative	494,368	98%
New negative	461,846	93%
New positive	29,949	6%
New exposed infants	1,047	0%
New inconclusive	1,526	0%
Confirmatory results (previous positive clients)	8,764	2%
Confirmatory positive	8,139	93%
Confirmatory inconclusive	625	7%

## HTC site report

Malawi (national)

2014 Q3 (1st month of quarter, 2nd month of quarter, 3rd month of quarter)

### HTC client details

\*

#### Partner / Family HTC referral slips

Sum of slips given	13,895	100%
Total clients presenting with referral slip	3,067	22%
Total failed referrals (slips not returned)	10,828	78%

### Clients tested in the community

#### HTC client details

\*

#### Total HTC clients served

Total HIV tested	25,177	100%
------------------	--------	------

#### Sex

Males tested	10,353	41%
Females tested	14,824	59%
Females non-pregnant	10,861	73%
Females pregnant	3,963	27%

#### Age

Children 0-14 yrs	3,845	15%
Children below 12 mths (Age group A)	37	1%
Children 18 mths - 14 yrs (Age group B)	3,808	99%
Adults 15+ years	21,332	85%
Young adults 15-24 years (Age group C)	10,229	48%
Older adults 25+ yrs (Age group D)	11,103	52%

#### HTC access type

PITC	5,525	22%
Family Referral Slip (FRS)	55	0%
Other (VCT, etc.) HTC access	19,597	78%

#### HTC first time / repeat

Never tested before	9,226	37%
Previously accessed HTC	15,951	63%
Last negative	15,760	99%
Last positive	159	1%
Last exposed infant	26	0%
Last inconclusive	6	0%

#### Counseling session type / Partner present

Counseled with partner / partner present	3,428	14%
Counseled alone / Partner not present	21,749	86%

#### Outcome summary (HIV test)

Single test negative	24,286	96%
Single test positive	188	1%
Test 1&2 negative	20	0%
Test 1&2 positive	637	3%
Test 1&2 discordant	46	0%

## HTC site report

Malawi (national)

2014 Q3 (1st month of quarter, 2nd month of quarter, 3rd month of quarter)

### HTC client details

\*

#### Final result given to client

Results among clients never tested / last negative	25,064	100%
New negative	24,456	98%
New positive	529	2%
New exposed infants	14	0%
New inconclusive	65	0%
Confirmatory results (previous positive clients)	113	0%
Confirmatory positive	104	92%
Confirmatory inconclusive	9	8%

#### Partner / Family HTC referral slips

Sum of slips given	149	100%
Total clients presenting with referral slip	55	37%
Total failed referrals (slips not returned)	94	63%

### Clients at stand-alone HTC sites

#### HTC client details

\*

#### Total HTC clients served

Total HIV tested	22,116	100%
------------------	--------	------

#### Sex

Males tested	11,852	54%
Females tested	10,264	46%
Females non-pregnant	9,359	91%
Females pregnant	905	9%

#### Age

Children 0-14 yrs	629	3%
Children below 12 mths (Age group A)	2	0%
Children 18 mths - 14 yrs (Age group B)	627	100%
Adults 15+ years	21,487	97%
Young adults 15-24 years (Age group C)	7,715	36%
Older adults 25+ yrs (Age group D)	13,772	64%

#### HTC access type

PITC	7,080	32%
Family Referral Slip (FRS)	22	0%
Other (VCT, etc.) HTC access	15,014	68%

#### HTC first time / repeat

Never tested before	7,107	32%
Previously accessed HTC	15,009	68%
Last negative	14,537	97%
Last positive	292	2%
Last exposed infant	173	1%
Last inconclusive	7	0%

#### Counseling session type / Partner present

Counseled with partner / partner present	2,046	9%
Counseled alone / Partner not present	20,070	91%

## HTC site report

Malawi (national)

2014 Q3 (1st month of quarter, 2nd month of quarter, 3rd month of quarter)

### HTC client details

\*

#### Outcome summary (HIV test)

Single test negative	20,072	91%
Single test positive	2	0%
Test 1&2 negative	12	0%
Test 1&2 positive	1,974	9%
Test 1&2 discordant	56	0%

#### Final result given to client

Results among clients never tested / last negative	21,993	99%
New negative	19,919	91%
New positive	1,869	8%
New exposed infants	1	0%
New inconclusive	204	1%
Confirmatory results (previous positive clients)	123	1%
Confirmatory positive	123	100%
Confirmatory inconclusive	0	0%

#### Partner / Family HTC referral slips

Sum of slips given	476	100%
Total clients presenting with referral slip	22	5%
Total failed referrals (slips not returned)	454	95%

## Blood safety

Malawi (national)

2014 Q3 (1st month of quarter, 2nd month of quarter, 3rd month of quarter)

### Infect. disease screening among potential donors

\*

#### HIV screening

HIV testing not done	1,437	17%
Tested for HIV	6,984	83%
HIV negative	6,548	94%
HIV positive	436	6%

#### Hepatitis B screening

HepB testing not done	1,516	18%
Tested for Hepatitis B	6,905	82%
HepB Negative	6,597	96%
HepB Positive	308	4%

#### Hepatitis C screening

HepC testing not done	6,215	74%
Tested for Hepatitis C	2,206	26%
HepC Negative	2,162	98%
HepC Positive	44	2%

#### Syphilis screening

Syphilis testing not done	1,533	18%
Tested for Syphilis	6,888	82%
Syphilis Negative	6,625	96%
Syphilis Positive	263	4%

#### Malaria screening

Malaria testing not done	4,105	49%
Tested for malaria	4,316	51%
Malaria Negative	4,041	94%
Malaria Positive	275	6%

#### Summary screening outcome

Not donated	2,774	33%
Donated	5,647	67%
Screened for at least HIV, HepB and syphilis	5,141	91%
Screened for HIV, HepB, HepC, Syphilis, Malaria	1,237	24%
Screened for HIV, HepB, Syphilis	3,904	76%
Screened for HIV, HepB	0	0%
Screened for HIV only	1	0%
Screened with any other combination of tests	505	9%

### Cross-matching report

\*

#### Blood group typing (for units and patients)

Total blood group typing done	18,338	100%
-------------------------------	--------	------

#### Blood units cross-matched (by source)

Total blood units cross-matched	12,793	100%
Total units from MBTS (estimated)	7,146	56%
Total units from replacement donors	5,647	44%

#### Blood units cross-matched by patient group

Units cross-matched for maternity	2,595	20%
Units cross-matched for paediatrics	3,587	28%
Units cross-matched for other ward	6,611	52%

## Blood safety

Malawi (national)

2014 Q3 (1st month of quarter, 2nd month of quarter, 3rd month of quarter)

### Cross-matching report

\*

#### Transfusion reactions

Units transfused without adverse events	12,716	99%
Units with suspected transfusion reactions	14	0%
Units with confirmed transfusion reactions	63	0%

# HIV exposed child follow-up

Malawi (national)

2014 Q3 (1st month of quarter, 2nd month of quarter, 3rd month of quarter)

## Age 2 months

### Age cohort outcomes

\*

#### Total children in birth cohort

Total children registered	7,236	100%
---------------------------	-------	------

#### CPT status

On CPT	6,545	90%
Not on CPT	691	10%

#### HIV status

Current HIV infection status unknown	4,552	63%
HIV infection not confirmed, not ART eligible	4,511	99%
HIV infection not confirmed, ART eligible (PSHD)	41	1%
Current HIV infection status known	2,684	37%
Confirmed not infected	2,618	98%
Confirmed infected (ART eligible)	66	2%

#### ART eligibility summary

Not eligible for ART	7,129	99%
ART eligible	107	1%
ART not initiated	68	64%
Initiated ART	39	36%

#### Primary follow-up outcome

Discharged uninfected	4	0%
Continue follow-up	6,625	93%
Started ART	39	1%
Defaulted	467	7%
Died	16	0%

#### Transfers between sites

Total not transferred out	7,151	99%
Transferred out	85	1%

## Age 12 months

### Age cohort outcomes

\*

#### Total children in birth cohort

Total children registered	8,184	100%
---------------------------	-------	------

#### CPT status

On CPT	5,612	69%
Not on CPT	2,572	31%

#### HIV status

Current HIV infection status unknown	4,993	61%
HIV infection not confirmed, not ART eligible	4,963	99%
HIV infection not confirmed, ART eligible (PSHD)	30	1%
Current HIV infection status known	3,191	39%
Confirmed not infected	3,023	95%
Confirmed infected (ART eligible)	168	5%

## HIV exposed child follow-up

Malawi (national)

2014 Q3 (1st month of quarter, 2nd month of quarter, 3rd month of quarter)

### Age cohort outcomes

\*

#### ART eligibility summary

Not eligible for ART	7,986	98%
ART eligible	198	2%
ART not initiated	37	19%
Initiated ART	161	81%

#### Primary follow-up outcome

Discharged uninfected	41	1%
Continue follow-up	5,550	70%
Started ART	161	2%
Defaulted	2,110	27%
Died	72	1%

#### Transfers between sites

Total not transferred out	7,934	97%
Transferred out	250	3%

### Age 24 months

#### Age cohort outcomes

\*

#### Total children in birth cohort

Total children registered	7,268	100%
---------------------------	-------	------

#### CPT status

On CPT	1,088	15%
Not on CPT	6,180	85%

#### HIV status

Current HIV infection status unknown	4,274	59%
HIV infection not confirmed, not ART eligible	4,253	100%
HIV infection not confirmed, ART eligible (PSHD)	21	0%
Current HIV infection status known	2,994	41%
Confirmed not infected	2,811	94%
Confirmed infected (ART eligible)	183	6%

#### ART eligibility summary

Not eligible for ART	7,064	97%
ART eligible	204	3%
ART not initiated	23	11%
Initiated ART	181	89%

#### Primary follow-up outcome

Discharged uninfected	2,641	38%
Continue follow-up	878	12%
Started ART	181	3%
Defaulted	3,252	46%
Died	88	1%

#### Transfers between sites

Total not transferred out	7,040	97%
Transferred out	228	3%

2014 Q3 (Quarter)

**Registration details**

\*

**HCC clinic registrations**

Total HCC registrations	17,554	100%
-------------------------	--------	------

**Registration type**

Patients enrolled first time	16,512	94%
Patients re-enrolled	48	0%
Patients transferred in	994	6%

**Sex**

Males (all ages)	8,332	47%
Females (all ages)	9,222	53%
Non-pregnant	9,192	100%
Pregnant	30	0%

**Age at registration**

Adults 15+ yrs	6,491	37%
Children 0-14 yrs	11,063	63%
Children 24 months - 14 years	546	5%
Children below 24 months (exposed children)	10,517	95%
Children 2 - below 24 months	3,851	37%
Infants below 2 months	6,666	63%

**Reason for HCC registration**

Exposed infants	10,613	60%
Confirmed infected patients (pre-ART)	6,941	40%

2014 Q3 (Cumulative)

**Registration details**

\*

**HCC clinic registrations**

Total HCC registrations	306,436	100%
-------------------------	---------	------

**Registration type**

Patients enrolled first time	296,311	97%
Patients re-enrolled	1,003	0%
Patients transferred in	9,122	3%

**Sex**

Males (all ages)	130,366	43%
Females (all ages)	176,070	57%
Non-pregnant	175,200	100%
Pregnant	870	0%

**Age at registration**

Adults 15+ yrs	161,479	53%
Children 0-14 yrs	144,957	47%
Children 24 months - 14 years	14,967	10%
Children below 24 months (exposed children)	129,990	90%
Children 2 - below 24 months	65,444	50%
Infants below 2 months	64,546	50%

**Reason for HCC registration**

Exposed infants	129,711	42%
Confirmed infected patients (pre-ART)	176,725	58%

**Pre-ART follow-up outcome**

\*

**Primary follow-up outcomes**

Total retained in pre-ART	38,741	23%
Started ART	86,847	51%
Defaulted	43,127	25%
Died	1,859	1%

**Transfers between sites**

Total not transferred out	170,548	97%
Transferred out	6,177	3%

## Antenatal Care

Malawi (national)

2014 Q3 (1st month of quarter, 2nd month of quarter, 3rd month of quarter)

### New ANC registrations in reporting period

\*

#### Women with first visit in reporting period

New women registered	164,321	100%
----------------------	---------	------

### ANC cohort analysis

\*

#### Total women completing ANC in the reporting period

Total women in booking cohort	153,514	100%
-------------------------------	---------	------

#### Visits per woman

Women with 1 visit	34,188	22%
Women with 2 visits	40,542	26%
Women with 3 visits	44,282	29%
Women with 4 visits	27,842	18%
Women with 5+ visits	6,660	4%

#### Trimester of first visit

Started ANC 0-12 wks	14,086	9%
Started ANC 13+ wks	139,371	91%

#### Pre-eclampsia

No pre-eclampsia	151,405	99%
Pre-eclampsia	2,052	1%

#### TTV doses

0-1 TTV doses	76,708	50%
2+ TTV doses	76,749	50%

#### SP tablets

0 SP doses	18,226	12%
1 SP dose (1 x 3 tabs)	43,212	28%
6+ SP tablets (2 x 3 tabs)	92,019	60%

#### FeFo tablets

0-119 FeFo tablets	117,906	77%
120+ FeFo tablets	35,551	23%

#### Albendazole (Deworming)

0 Albend. doses	27,786	18%
1 Albend. dose	126,925	82%

#### ITN (bednets)

No ITN	44,153	28%
ITN received	111,037	72%

#### Syphilis status

Not tested for syphilis	141,766	92%
Tested for syphilis	11,691	8%
Syphilis negative	11,434	98%
Syphilis positive	257	2%

## Antenatal Care

Malawi (national)

2014 Q3 (1st month of quarter, 2nd month of quarter, 3rd month of quarter)

### ANC cohort analysis

\*

#### HIV status ascertainment

HIV status not ascertained	17,499	11%
HIV status ascertained	135,958	89%
Valid previous test result	9,317	7%
Previous negative	4,645	50%
Previous positive	4,672	50%
New test at ANC	126,641	93%
New negative	121,017	96%
New positive	5,624	4%

#### HIV status summary

Total women HIV negative	125,662	92%
Total women HIV positive	10,296	8%

#### CPT status (among HIV pos)

Not on CPT	779	8%
On CPT	9,517	92%

#### Final PMTCT regimen mother

No ARVs	744	7%
Any ARVs	9,550	93%
ART (by time of initiation)	9,550	100%
Already on ART when starting ANC	4,241	44%
Started ART at 0-27 weeks of pregnancy	4,126	43%
Started ART at 28+ weeks of preg.	1,183	12%

#### Baby's ARVs dispensed

No ARVs dispensed for infant	2,260	22%
ARVs dispensed for infant	8,036	78%

# Maternity

Malawi (national)

2014 Q3 (1st month of quarter, 2nd month of quarter, 3rd month of quarter)

## Maternal details

\*

### Admissions in the reporting period

Total admissions (referrals double-counted)	137,343	100%
Not referred to other site (total women)	129,857	95%
Referred out before delivery (multiple admissions)	7,486	5%

### HIV status ascertainment

HIV status not ascertained	6,498	5%
HIV status ascertained	130,167	95%
Valid previous test result	127,052	98%
Previous negative	117,438	92%
Previous positive	9,614	8%
New test at maternity	3,115	2%
New negative	2,836	91%
New positive	279	9%

### HIV status summary

Total women HIV negative	120,274	92%
Total women HIV positive	9,893	8%

### ARVs during pregnancy (among HIV pos)

No ARV in pregnancy	258	3%
Any ARVs	9,635	97%
ART (by time of initiation)	9,635	100%
ART initiated before pregnancy	6,579	68%
ART initiated in 1st / 2nd trimester	1,602	17%
ART initiated in 3rd trimester	1,190	12%
ART initiated during labour	264	3%

### Obstetric complications

No obstetric complications	120,176	88%
Any obstetric complications	16,489	12%
Haemorrhage	2,775	17%
Haemorrhage ante-partum	843	30%
Haemorrhage post-partum	1,932	70%
Obstr / prol labour	5,592	34%
(pre-) Eclampsia	1,348	8%
Maternal sepsis	141	1%
Ruptured uterus	145	1%
Other obstetric complications	6,488	39%

### Emergency obstetric care

Oxytocin	125,613	95%
Anticonvulsive	653	0%
Antibiotics	4,850	4%
Blood transfusion	462	0%
Manual removal of placenta	243	0%

### Vitamin A

Vit A not given	46,445	34%
Vit A given	90,220	66%

# Maternity

Malawi (national)

2014 Q3 (1st month of quarter, 2nd month of quarter, 3rd month of quarter)

## Maternal details

\*

### Staff conducting delivery

Category A: MO, CO, nurse/midwife, MA	123,481	96%
Category B: PA, WA, HSA	844	1%
Category C: Other	4,854	4%

### Mother survival

Mother alive	129,068	100%
Mother died	111	0%

## Infant details

\*

### Single babies / multiple deliveries

Total babies delivered	131,616	100%
Single babies	127,129	97%
Twin / multiple babies	4,487	3%

### Delivery place

Total deliveries at a health facility	126,611	96%
This facility	126,239	100%
Other facility	372	0%
Total deliveries before reaching the facility	5,005	4%
In transit	3,216	64%
Home / TBA	1,789	36%

### Delivery mode

Spontaneous vaginal	118,375	90%
Vacuum extraction	1,780	1%
Breech	2,267	2%
Caesarean section	9,194	7%

### Infant complications

No infant complications	115,788	88%
Total infants with complications	15,828	12%
Prematurity	3,581	23%
Weight less 2500g	4,949	31%
Asphyxia	4,877	31%
Sepsis	607	4%
Other newborn complication	1,814	11%

### Infant survival

Total live births	129,284	98%
Discharged alive	128,029	99%
Neonatal deaths	1,255	1%
Stillbirths	2,332	2%
Stillbirth, fresh	1,214	52%
Stillbirth, macerated	1,118	48%

## Maternity

Malawi (national)

2014 Q3 (1st month of quarter, 2nd month of quarter, 3rd month of quarter)

### Infant details

\*

#### HIV exposure / ARV proph. (among discharged alive)

Infants with unknown HIV exposure status	5,368	4%
Infants with known HIV exposure status	122,661	96%
Not HIV exposed	113,574	93%
HIV exposed	9,087	7%
Received no ARVs	667	7%
Received ARVs	8,420	93%
Nevirapine	8,420	100%

#### Breastfeeding initiated

BF not started within 60min	11,651	9%
BF started within 60min	119,965	91%

#### Tetracycline eye ointment given

TO not given	18,208	14%
TO given	113,408	86%

# ART cohort analysis

Malawi (national)

2014 Q3 (Quarter)

## Registration details

\*

### ART clinic registrations

Total ART clinic registrations	37,082	100%
--------------------------------	--------	------

### Registration type

First time ART initiations (total patients)	29,893	81%
ART re-initiations	435	1%
ART transfers in	6,754	18%

### Sex

Males	13,046	35%
Females	24,036	65%
Non-pregnant	17,032	71%
Pregnant	7,004	29%

### Age at ART initiation

Adults 15+ yrs	34,202	92%
Children 0-14 yrs	2,880	8%
Children 2-14 yrs	2,204	77%
Children below 24 mths	676	23%

### Reason for starting ART

Presumed severe HIV Disease	113	0%
Confirmed HIV infection	36,969	100%
WHO stage 1 or 2	25,101	68%
Total lymphocytes <threshold	5	0%
CD4 below threshold	15,503	62%
CD4 unknown or >threshold	9,593	38%
PCR infants	123	1%
Children 12-59 mths	686	7%
Pregnant women	6,974	73%
Breastfeeding mothers	1,810	19%
WHO stage 3	9,988	27%
WHO stage 4	1,626	4%
Unknown / reason outside of guidelines	254	1%

### TB at ART initiation

Never TB / TB > 24 months ago	35,952	97%
TB within the last 24 months	411	1%
Current episode of TB	719	2%

### Kaposi's sarcoma at ART initiation

No KS	36,772	99%
Patients with KS	310	1%

# ART cohort analysis

Malawi (national)

2014 Q3 (Cumulative)

## Registration details

\*

### ART clinic registrations

Total ART clinic registrations	929,253	100%
--------------------------------	---------	------

### Registration type

First time ART initiations (total patients)	745,133	80%
ART re-initiations	9,709	1%
ART transfers in	174,411	19%

### Sex

Males	334,355	36%
Females	594,898	64%
Non-pregnant	492,131	83%
Pregnant	102,767	17%

### Age at ART initiation

Adults 15+ yrs	848,820	91%
Children 0-14 yrs	80,433	9%
Children 2-14 yrs	62,130	77%
Children below 24 mths	18,303	23%

### Reason for starting ART

Presumed severe HIV Disease	3,337	0%
Confirmed HIV infection	925,916	100%
WHO stage 1 or 2	386,122	42%
Total lymphocytes <threshold	243	0%
CD4 below threshold	261,573	68%
CD4 unknown or >threshold	124,306	32%
PCR infants	2,634	2%
Children 12-59 mths	5,021	4%
Pregnant women	83,837	67%
Breastfeeding mothers	32,814	26%
WHO stage 3	439,087	47%
WHO stage 4	95,757	10%
Unknown / reason outside of guidelines	4,950	1%

### TB at ART initiation

Never TB / TB > 24 months ago	860,023	93%
TB within the last 24 months	35,649	4%
Current episode of TB	33,581	4%

### Kaposi's sarcoma at ART initiation

No KS	910,672	98%
Patients with KS	18,581	2%

# ART cohort analysis

Malawi (national)

2014 Q3 (Cumulative)

## ART outcomes

\*

### Primary follow-up outcomes

Total alive on ART	521,319	69%
Alive on ART at site of last registration	519,311	100%
ART patients in transit between sites	2,008	0%
Defaulted	158,580	21%
Stopped ART	3,418	0%
Total died	71,525	9%
Died month 1	18,337	26%
Died month 2	11,522	16%
Died month 3	6,721	9%
Died month 4+	34,945	49%

### Transfers between sites

Total not transferred out	752,834	81%
Transferred out	176,419	19%

### ART regimens

First line regimens	513,600	99%
Adult formulation	488,020	95%
Regimen 0A	80	0%
Regimen 1A	1,925	0%
Regimen 2A	26,686	5%
Regimen 3A	158	0%
Regimen 4A	685	0%
Regimen 5A	454,763	93%
Regimen 6A	3,723	1%
Paed. formulation	25,580	5%
Regimen 0P	123	0%
Regimen 1P	561	2%
Regimen 2P	24,401	95%
Regimen 3P	109	0%
Regimen 4P	386	2%
Second line regimens	5,240	1%
Adult formulation	4,582	87%
Regimen 7A	3,885	85%
Regimen 8A	697	15%
Paed. Formulation	658	13%
Regimen 9P	658	100%
Other regimen (adult / paed)	471	0%

### Adherence

Adherence unknown (not recorded)	5,242	1%
Adherence recorded	514,069	99%
0-3 doses missed	469,725	91%
4+ doses missed	44,344	9%

### ART side effects

Side effects unknown (not recorded)	4,019	1%
Side effects recorded	515,292	99%
No side effects	507,137	98%
Any side effects	8,155	2%

# ART cohort analysis

Malawi (national)

2014 Q3 (Cumulative)

## ART outcomes

\*

### Current TB status among ART patients (ICF)

ICF not done (Current TB status unknown/ not circ)	10,712	2%
ICF done	508,599	98%
TB not suspected	501,349	99%
TB suspected	5,583	1%
TB confirmed	1,667	0%
TB confirmed, not on treatment	139	8%
TB confirmed, on TB treatment	1,528	92%

2014 Q3 (Quarter)

**12 month survival children****Survival and retention in ART program**

\*

**ART cohort registration group outcomes**

Total ART clinic registrations	3,181	100%
Transfers out (double counted)	346	11%
Total not transferred out (patients in cohort)	2,835	89%
Total alive on ART	2,213	78%
Total not retained	622	22%
Defaulted	487	78%
Stopped ART	10	2%
Died	125	20%

**12 month survival all ages****Survival and retention in ART program**

\*

**ART cohort registration group outcomes**

Total ART clinic registrations	33,776	100%
Transfers out (double counted)	3,582	11%
Total not transferred out (patients in cohort)	30,194	89%
Total alive on ART	23,585	78%
Total not retained	6,609	22%
Defaulted	5,403	82%
Stopped ART	67	1%
Died	1,139	17%

**24 month survival all ages****Survival and retention in ART program**

\*

**ART cohort registration group outcomes**

Total ART clinic registrations	39,356	100%
Transfers out (double counted)	5,351	14%
Total not transferred out (patients in cohort)	34,005	86%
Total alive on ART	24,740	73%
Total not retained	9,265	27%
Defaulted	7,253	78%
Stopped ART	125	1%
Died	1,887	20%

**36 month survival all ages****Survival and retention in ART program**

\*

**ART cohort registration group outcomes**

Total ART clinic registrations	33,916	100%
Transfers out (double counted)	6,516	19%
Total not transferred out (patients in cohort)	27,400	81%
Total alive on ART	19,844	72%
Total not retained	7,556	28%
Defaulted	5,352	71%
Stopped ART	146	2%
Died	2,058	27%

# ART survival analysis

Malawi (national)

2014 Q3 (Quarter)

## 48 month survival all ages

### Survival and retention in ART program

\*

#### ART cohort registration group outcomes

Total ART clinic registrations	23,096	100%
Transfers out (double counted)	5,745	25%
Total not transferred out (patients in cohort)	17,351	75%
Total alive on ART	11,099	64%
Total not retained	6,252	36%
Defaulted	4,221	68%
Stopped ART	88	1%
Died	1,943	31%

## 60 month survival all ages

### Survival and retention in ART program

\*

#### ART cohort registration group outcomes

Total ART clinic registrations	21,114	100%
Transfers out (double counted)	5,897	28%
Total not transferred out (patients in cohort)	15,217	72%
Total alive on ART	9,050	59%
Total not retained	6,167	41%
Defaulted	4,134	67%
Stopped ART	101	2%
Died	1,932	31%

## 72 month survival all ages

### Survival and retention in ART program

\*

#### ART cohort registration group outcomes

Total ART clinic registrations	20,211	100%
Transfers out (double counted)	5,619	28%
Total not transferred out (patients in cohort)	14,592	72%
Total alive on ART	8,384	57%
Total not retained	6,208	43%
Defaulted	3,875	62%
Stopped ART	108	2%
Died	2,225	36%

## 84 month survival all ages

### Survival and retention in ART program

\*

#### ART cohort registration group outcomes

Total ART clinic registrations	15,816	100%
Transfers out (double counted)	4,742	30%
Total not transferred out (patients in cohort)	11,074	70%
Total alive on ART	5,748	52%
Total not retained	5,326	48%
Defaulted	3,261	61%
Stopped ART	86	2%
Died	1,979	37%

# ART survival analysis

Malawi (national)

2014 Q3 (Quarter)

## 96 month survival all ages

### Survival and retention in ART program

\*

#### ART cohort registration group outcomes

Total ART clinic registrations	12,162	100%
Transfers out (double counted)	3,763	31%
Total not transferred out (patients in cohort)	8,399	69%
Total alive on ART	4,113	49%
Total not retained	4,286	51%
Defaulted	2,130	50%
Stopped ART	66	2%
Died	2,090	49%

## 108 month survival all ages

### Survival and retention in ART program

\*

#### ART cohort registration group outcomes

Total ART clinic registrations	7,554	100%
Transfers out (double counted)	2,278	30%
Total not transferred out (patients in cohort)	5,276	70%
Total alive on ART	2,285	43%
Total not retained	2,991	57%
Defaulted	1,278	43%
Stopped ART	44	1%
Died	1,669	56%

## 120 month survival all ages

### Survival and retention in ART program

\*

#### ART cohort registration group outcomes

Total ART clinic registrations	2,821	100%
Transfers out (double counted)	859	30%
Total not transferred out (patients in cohort)	1,962	70%
Total alive on ART	869	44%
Total not retained	1,093	56%
Defaulted	496	45%
Stopped ART	28	3%
Died	569	52%

## 6 month survival OptionB+

### Survival and retention in ART program

\*

#### ART cohort registration group outcomes

Total ART clinic registrations	8,164	100%
Transfers out (double counted)	509	6%
Total not transferred out (patients in cohort)	7,655	94%
Total alive on ART	5,992	78%
Total not retained	1,663	22%
Defaulted	1,603	96%
Stopped ART	14	1%
Died	46	3%

2014 Q3 (Quarter)

**12 month survival OptionB+****Survival and retention in ART program**

\*

**ART cohort registration group outcomes**

Total ART clinic registrations	8,318	100%
Transfers out (double counted)	712	9%
Total not transferred out (patients in cohort)	7,606	91%
Total alive on ART	5,544	73%
Total not retained	2,062	27%
Defaulted	1,976	96%
Stopped ART	23	1%
Died	63	3%

**24 month survival OptionB+****Survival and retention in ART program**

\*

**ART cohort registration group outcomes**

Total ART clinic registrations	9,852	100%
Transfers out (double counted)	1,073	11%
Total not transferred out (patients in cohort)	8,779	89%
Total alive on ART	6,058	69%
Total not retained	2,721	31%
Defaulted	2,541	93%
Stopped ART	61	2%
Died	119	4%

**36 month survival OptionB+****Survival and retention in ART program**

\*

**ART cohort registration group outcomes**

Total ART clinic registrations	6,203	100%
Transfers out (double counted)	1,009	16%
Total not transferred out (patients in cohort)	5,194	84%
Total alive on ART	3,490	67%
Total not retained	1,704	33%
Defaulted	1,497	88%
Stopped ART	56	3%
Died	151	9%

# STI site report

Malawi (national)

2014 Q3 (1st month of quarter, 2nd month of quarter, 3rd month of quarter)

## STI clients treated in the reporting period

\*

### Total STI clients

Total STI clients treated	54,427	100%
Index patients treated (symptomatic)	43,010	79%
Partners treated	11,417	21%

### Sex

Males	22,180	41%
Females	32,247	59%
Non-pregnant	28,085	87%
Pregnant	4,162	13%

### Age group

Age group A (0-19 years)	4,363	8%
Age group B (20-24 years)	13,050	24%
Age group C (25+ years)	37,014	68%

### Client type

Symptomatic cases	47,547	87%
Index cases	43,010	90%
Partners symptomatic	4,537	10%
Partners asymptomatic	6,880	13%

### STI treatment history

Never treated for STI	40,330	74%
Previously treated for STI	14,097	26%
Old >3 months ago	9,006	64%
Recent ≤3 months ago	5,091	36%

### STI syndromic diagnosis

GUD	9,770	17%
UD	13,474	23%
AVD	17,143	29%
Low risk	6,971	41%
High risk	10,172	59%
LAP	9,611	16%
SS	953	2%
BU	655	1%
BA	1,180	2%
NC	263	0%
Genital Warts	583	1%
Syphilis RPR VDRL	1,197	2%
Other STI	3,733	6%

### STI partner notification

Total partner notification slips issued	12,507	100%
Total partners returned	11,417	91%
Total partners not seen	1,090	9%

## STI site report

Malawi (national)

2014 Q3 (1st month of quarter, 2nd month of quarter, 3rd month of quarter)

### STI clients treated in the reporting period

\*

#### HIV test / ART status

HIV status not ascertained	27,308	50%
HIV status ascertained	27,119	50%
HIV negative (new test)	19,944	74%
HIV positive	7,175	26%
New positive	1,495	21%
Previous positive	5,680	79%
Not on ART	1,484	26%
On ART	4,196	74%

#### STI clients referred for services

Lab	548	3%
Gynae review	380	2%
Surgical review	438	2%
Repeat HTC	15,199	79%
ART (for assessment)	880	5%
PMTCT	130	1%
Other (service referrals)	1,628	8%