Government of Malawi Ministry of Health

Integrated HIV Program Report
January -March 2015

- *Integrated HIV Program Supervision*
- *HIV Testing and Counselling / Early Infant Diagnosis*
- *Blood Safety*
- *Post Exposure Prophylaxis*
- *HIV Exposed Child Follow-Up*
- *Pre-ART*
- *Prevention of Mother to Child Transmission / Antiretroviral Therapy*
- *TB / HIV*
- *Sexually Transmitted Infections*
- *Supply of HIV Program Commodities*

All current and past issues of our program reports and many other resources are published on the HIV Department website: ***www.hiv.health.gov.mw***

1 EXECUTIVE SUMMARY 2
2 INTEGRATED HIV PROGRAM OVERVIEW 3
3 SUPPORTIVE SITE SUPERVISION 4
3.1 Methods 4
3.2 Supervision Outcomes 4
4 INVENTORY OF SITES AND SERVICES 5
4.1 Sites and Services 5
4.2 Staffing of HIV Services 6
5 HIV TESTING AND COUNSELLING PROGRAM OUTPUTS 7
5.1 HTC data 7
5.2 HTC access type 8
5.3 Age and sex distribution among HTC clients 8
5.4 First time, repeat and confirmatory test results 8
6 DNA-PCR TESTING FOR EARLY INFANT DIAGNOSIS OF HIV (EID) 9
7 BLOOD SAFETY 10
8 POST EXPOSURE PROPHYLAXIS (PEP) 11
9 PROVIDER-INITIATED FAMILY PLANNING (PIFP) 11
10 COTRIMOXAZOLE PREVENTIVE THERAPY (CPT) 11
11 TB / HIV INTERVENTIONS 12
11.1 Intensified Case Finding (ICF) 12
11.2 Isoniazid Preventive Therapy (IPT) 12
12 HIV-RELATED DISEASES 13
13 HIV-EXPOSED CHILD FOLLOW-UP 13
13.1 Methods and Definition of Indicators 13
13.2 HIV Exposed Child Registration Data 14
13.3 Birth Cohort Outcomes 14
14 PRE-ART 15
14.1 Pre-ART Registration Data 15
14.2 Cumulative Pre-ART Follow-up Outcomes 15
15 PMTCT / ART 15
15.1 Data Sources and Reporting Methods 15
15.2 ARV Coverage among Pregnant / Breastfeeding Women and Exposed Infants 16
15.3 HIV Services at ANC 17
15.4 HIV Services at Maternity 18
16 ART ACCESS AND FOLLOW-UP OUTCOMES 19
16.1 New ART Registrations during Q1 2015 19
16.2 Cumulative ART Registrations up March 2015 20
16.3 ART Outcomes 21
16.4 ART Cohort Survival Analysis 23
16.5 Viral Load (VL) Monitoring 27
17 TB / HIV MANAGEMENT 28
18 STI TREATMENT 28
18.1 Access to STI treatment and coverage 29
18.2 Client Type and STI History 29
18.3 HIV Status 29
18.4 STI Syndromes 29
18.5 Referrals 29
19 SUPPLY OF HIV PROGRAM COMMODITIES 30
19.1 Quantification and procurement planning 30
19.2 Quarterly distribution of HIV Commodities 30
19.3 Quarterly logistics support during Q1 2015 ART/PMTCT supervision 30
19.4 National Stock Status of HIV Commodities 31
20 TRAINING AND MENTORING 33
21 PARTICIPANTS IN Q1 2015 SUPERVISION (SITE VISITS 13 APRIL-1 MAY 2015) 34
22 APPENDIX (FULL NATIONAL HIV PROGRAM DATA) 34

1 Executive Summary

This is the 15th quarterly HIV Program report after implementation of the 2011 Integrated Clinical HIV Guidelines in July 2011. A summary of the key achievements between **January and March 2015** is provided below:

- Scale-up of integrated HIV services had reached the following number of sites:
  - **724** static (579 within and 145 outside of health facilities) and 188 outreach HTC sites
  - **714** (static) ART sites
  - **620** PMTCT sites (Option B+, all included in ART sites above)
  - **657** Pre-ART sites
  - **641** sites with HIV-exposed child follow-up
- **500,359** persons were tested and counselled for HIV; **157,080 (31%)** accessed HTC for the first time; **343,279 (69%)** were repeat testers and **9,992 (3%)** of these received confirmatory testing (after having tested positive in the past). This is equivalent to **40%** confirmatory testing coverage among 25,270 patients initiating ART this quarter. **31,491 (6%)** clients received a positive result for the first time.
- **18,765 (97%)** of 19,436 blood units collected were screened for (at least) HIV, hepatitis B and syphilis.
- **148,838 (91%)** of 164,275 women at ANC had their HIV status ascertained; **11,727 (8%)** of these were HIV positive. **120,651 (96%)** of 129,368 women at maternity had their HIV status ascertained **9,142 (8%)** of these were HIV positive.
- **25,270** patients started ART this quarter.
- **551,566** patients were alive and on ART by end of March 2015. This means that **55%** of the estimated 1 million HIV positive population was on ART. [1] Estimated ART coverage among people in need for treatment[1] was **48%** (47,962 / 101,000) for children (<15 years) and **65%** (503,604 / 779,000) for adults.
- **76%** of adults and **77%** of children were retained alive on ART at 12 months after initiation. Actual retention rates are thought to be about **10%** higher due to misclassification of 'silent transfers' as defaulters in clinic-based survival/retention analysis. (see section 16.4)
- **481,309 (93%)** of 517,067 patients on first line adult ART were on regimen 5A (TDF/3TC/EFV).
- **11,583** [2] **(85%)** of an estimated **13,317** [1] HIV infected pregnant women in Malawi were on ART this quarter. **6,266 (59%)** of these were already on ART when getting pregnant and **4,317 (41%)** started ART during pregnancy/delivery.
- An additional **1,584** [2] breastfeeding women started ART due to ***Option B+*** (in WHO stage 1/2)
- **77%, 72%, 69%** and **70%** of women started under ***Option B+*** were retained on ART at **6, 12, 24** and **36 months** after initiation, respectively.
- **8,499 (7%)** of infants discharged alive from maternity were known to be HIV exposed, **7,866 (93%)** of these received ARV prophylaxis (nevirapine). **6,750 (65%)** were enrolled in exposed child follow-up before age 2 months.
- A total of **10,452** HIV exposed children and **6,223** pre-ART patients were enrolled for follow-up in *HIV Care Clinics (HCC)* during this quarter.

[1] 2015 Spectrum estimates based on 2014 definition of eligibility for ART in Malawi (CD4<500, Option B+, UT for U5).

[2] Adjusted for double counting due to patient transfers / 'failed ART initiations' among women lost to follow-up within 6 months of ART registration.

2 Integrated HIV Program Overview

Malawi implemented a revised HIV Program in all health facilities following the release of the ***2011 Malawi Integrated Clinical HIV Guidelines***. The second edition of these guidelines was published in March 2014 and implementation of revised policies commenced in April 2014. Key policies include:

- **PMTCT Option B+:** Universal life-long ART for all HIV infected pregnant and breastfeeding women regardless of clinical or immunological stage.

- Standard **HIV exposed child follow-up** to age 24 months. This program aims to improve early infant diagnosis and ART initiation using DNA-PCR testing for all infants from age 6 weeks; rapid antibody testing is considered diagnostic from age 12 months and repeated at 24 months. HIV exposed child enrolment and follow-up should be integrated with maternal ART follow-up (Option B+) to improve retention and adherence.

- **Early ART initiation**: universal ART for children under 5 years (confirmed HIV infection, CD4% no longer required), children over 5 years and adults with a **CD4** count **≤500**, patients with HIV and hepatitis B co-infection.

- Transition to a **new first line ART regimens** for adults (Malawi regimen 5A: tenofovir / lamivudine / efavirenz) and children (Malawi regimen 2: zidovudine / lamivudine / nevirapine), including provision of paediatric ARV formulations for all children under 25kg. Transition to 5A was completed by end 2013.

- Standardized **pre-ART services** for all HIV-infected persons not yet eligible for ART. The pre-ART program aims to reduce the incidence of HIV-related diseases and to enable early ART initiation based on the new CD4 cell threshold (500) through scheduled CD4 count monitoring.

- Provider-initiated provision of **contraceptives and condoms** for all adults in pre-ART and ART clinics to reduce the rate of unwanted pregnancies among HIV-infected adults and to reduce HIV-transmission between sexual partners.

- Isoniazid preventive therapy (**IPT**) for pre-ART patients to reduce the incidence of TB and intensified TB case finding (**ICF**) for all patients in pre-ART and ART follow-up to enable early diagnosis and treatment of TB and to reduce TB transmission in HIV clinics.

- Roll-out of scheduled **viral load monitoring** to improve early detection of treatment failure and initiation of second line ART.

Implementation of PMTCT Option B+ requires provision of ART services at all health facilities with Maternal and Child Health services. This required a massive acceleration of the **decentralization of ART services** from 303 (static) sites in June 2011 to currently 714 sites.

3 Supportive Site Supervision

3.1 Methods

The Department for HIV and AIDS has coordinated quarterly supportive supervision visits to all health facilities with ART services since the start of the national treatment program in 2004. Supervision teams are composed of: experienced HIV clinicians; nurses and M&E staff from health facilities in the public and private sector; district and zonal PMTCT and ART coordinators; program officers and technical staff from the Department for HIV and AIDS; technical staff from implementing partners. The TB and HIV programs are working towards a full integration of their respective site supervision exercises.

Each quarter, a one day pre-supervision meeting is organised for all supervisors participating in the upcoming round to share program updates, discuss observations from the previous round, distribute materials and organise logistics, transport and accommodation.

Standard supervision forms are used to guide implementation of the supervision protocol, to update site information and collect M&E reports. Custom forms with previous data for each site are printed from the HIV Department database. The supervision forms include:

- Contact details of HIV service providers at each site
- Quality of service checklist
- Follow up on action points noted during the previous visit
- Next visit date
- M&E reports from HTC, ANC, maternity, exposed child and pre-ART follow-up, ART and TB
- Physical drug stock-level assessment
- Identification of sites in urgent need of clinical mentoring
- Semi-structured feedback and performance rating for the supervision teams by facility staff

One copy of the supervision form is returned to the Department for HIV and AIDS, where data are entered in a custom MS Access database to produce national reports and to manage program logistics and the commodity supply chain. A second copy of the supervision form is left at the sites.

The supervision protocol includes a systematic review and verification of primary records (patient cards and registers) at all sites. This effectively provides a quarterly quality audit for M&E records, which has resulted in exceptional accuracy and completeness of HIV Program data in Malawi. At the same time, the systematic chart review helps to identify complex cases or deviations from clinical protocol, allowing the supervision team to provide targeted mentoring and clinical advice. The quarterly supervision exercise also aims to boost staff morale and motivation through *Certificates of Excellence* that are awarded by MOH to sites with an excellent score on the quality of service checklist. A growing number of health workers from sites all over the country participate as supervisors in this quarterly exercise and this has strengthened the national HIV Program identity and has greatly facilitated communication between program staff at the national, zonal, district and facility level.

The HTC Program usually conducts a separate supportive site supervision exercise each quarter, targeting a sample of HTC sites both within and outside of health facilities. Supervision teams consist of district, zonal and national level HTC coordinators, supported by implementing partners.

3.2 Supervision Outcomes

712 public and private sector facilities were visited for **clinical HIV program supervision** between 13th April and 1st May 2015.

The large number of sites was covered by **95** supervisors working in **23** teams that spent a total of **2,024 working hours** at the sites. Each site visit lasted on average **2.8** hours, but up to 2 days were spent at the busiest sites. **345 (48%)** sites were awarded a *Certificate of Excellence* for **excellent performance.** The

number of sites with excellent performance is similar higher than previous quarter 321. **51 (7%)** sites had significant weaknesses and were rated to require **intensive mentoring**. The capacity to provide site mentoring will need to be further expanded.

Table 1: Outcomes of integrated HIV services supervision for 2015 Q1

Zone	Total facil. visited*	Supervision hours spent at facilities		Performance (# and % of sites)	
		Total	Average per site	Excellent perform.	Mentoring needed
NZ	122	320	2.7	63 52%	8 7%
CEZ	99	254	2.6	47 47%	10 10%
CWZ	160	449	2.9	73 46%	6 4%
SEZ	166	503	3	85 51%	13 8%
SWZ	167	498	3	77 46%	14 8%
Malawi	714	2,024	2.8	345 48%	51 7%

* includes facilities that were visited for assessment of readiness, but that may have not (yet) been designated to provide integrated HIV services.

Table 1 provides a summary of the supervision outcomes by zone. Most facilities were using the standard national M&E tools. **133** sites had cumulatively registered more than 2,000 ART patient and **43** of these had registered more than 5,000. **51 (38%)** of these high burden sites were using electronic data system for ART (EDS). Some NGO supported sites were using custom tools compatible with the national standard reporting requirements.

4 Inventory of Sites and Services

4.1 Sites and Services

There were **724** static and **188** outreach HTC sites in Q1 2015; **145** of these were outside of health facilities.

Table 2: Facilities with integrated HIV services in the 5 Zones. Availability of services defined by performance (at least 1 patient enrolled) during 2015 Q1

Zone	Total fac.(1)	Facilities providing HIV services				CD4 count machines (2)		
		Exp. child	Pre-ART	PMTCT B+	ART	Installed	Functional	Results
NZ	130	109 84%	108 83%	96 74%	118 91%	25 19%	20 80%	2,412
CEZ	99	93 94%	92 93%	90 91%	97 98%	17 17%	15 88%	1,405
CWZ	162	132 81%	134 83%	135 83%	160 99%	27 17%	24 89%	3,467
SWZ	168	146 87%	163 97%	139 83%	165 98%	41 24%	33 80%	8,183
SEZ	165	160 97%	159 96%	159 96%	164 99%	51 31%	47 92%	5,510
Malawi	724	640 88%	656 91%	619 85%	714 99%	161 22%	139 86%	20,977

(1) Total facilities in the public / private sector designated to provide integrated HIV services in this quarter. Individual site selection is reviewed and may change each quarter.

(2) CD4 machines that have produced at least 1 result during the reporting period are defined as functional.

Table 2 shows the distribution of the **723** sites designated to provide clinical HIV services in Q1 2015, by zone. At the national level, there were **714** (static) sites with at least one patient on ART, **619** sites had enrolled women under PMTCT Option B+; **656** sites were providing pre-ART services. **640** had enrolled HIV exposed children for follow-up. ART services were now available at

almost all designated sites in the 5 zones. The SEZ and CWZ had reached 99% of designated sites with ART services. SEZ had reached 96% of designated sites with Option B+.

CD4 count machines (including 'point of care' machines) were installed at **161** sites, and **139** (86%) of these had produced at least 1 result during Q1 2015. The total number of CD4 results produced decreased from 24,485 in Q4 2014 to **20,977** during Q1 2015. 39% of these outputs were generated by 33 machines in the SW zone, implying that many CD4 machines continued to experience down-time or to be running considerably below capacity. The raised CD4 count threshold for ART eligibility may have also resulted in a decrease in the number of pre-ART patients requiring CD4 count monitoring as a large proportion will now be started on ART after an initial CD4 test.

4.2 Staffing of HIV Services

4.2.1 HTC Services

The Department for HIV and AIDS has maintained a dedicated system for professional registration and performance tracking for HTC Providers since 2011. This separate registration system is needed because HTC providers include lay persons with HTC training who are not registered with any other professional body. All HTC providers are issued with a unique ID and a professional logbook for documentation of duty stations, HTC trainings, sit-in observation and proficiency testing results. Logbook holders are requested to record the total number of tests done at the end of each month. Logbooks are routinely reviewed during quarterly supervision and key performance data for each provider are summarized on the site supervision forms.

	2014 Q2		2014 Q3		2014 Q4		2015 Q1	
Sites visited	**697**		**720**		**718**		**714**	
Sites with any tests done	652	*94%*	668	*93%*	660	*92%*	663	*93%*
Sites with registered HTC staff	645	*93%*	657	*91%*	659	*92%*	667	*93%*
Total HTC staff at visited sites	**3,445**		**3,583**		**3,574**		**3,673**	
Staff with any test done	2,064	*60%*	2,276	*64%*	2,195	*61%*	2,234	*61%*
Staff with 300+ tests done this quarter	255	*11%*	322	*13%*	286	*11%*	268	*11%*
Logbooks reviewed	2,335	*68%*	2,554	*71%*	2,526	*71%*	2,541	*69%*
HTC staff participating in PT this quarter	14	*1%*	1,380	*54%*	170	*7%*	1,641	*65%*
Total tests (HTC register)	**435,976**		**550,436**		**464,292**		**500,086**	
Tests accounted for by individual staff	299,233	*69%*	355,216	*65%*	331,981	*72%*	321,221	*64%*
Source: logbooks	278,053	*93%*	330,271	*93%*	303,046	*65%*	290,888	*91%*
Source: HTC register	21,180	*7%*	24,945	*7%*	28,935	*10%*	30,333	*9%*
Total tests by staff with 300+ tests	129,856	*43%*	154,494	*43%*	145,300	*44%*	134,272	*42%*

667 (93%) of the 714 visited facilities had registered HTC providers and **663** (93%) sites had performed at least one test during Q1 2015. **2,541 (69%)** of **3,673** HTC providers had their logbooks available for review.

According to the 2,541 reviewed logbooks, **1,641 (65%)** HTC providers had participated in proficiency (panel) testing (PT) this quarter. This exceeds the participation rates from the previous 3 quarters, but it implies that only about 2/3 of HTC providers underwent PT this quarter. Only 69% of providers at the visited sites had their logbooks available for review and documentation of PT may have been incomplete. The national HIV reference laboratory is organizing six monthly PT rounds for all practising HTC providers. PT panels are distributed by the district lab supervisors who may not be able to reach all sites in their districts.

321,221 (64%) of all 500,359 tests conducted this quarter (according to HTC register reports) were accounted for by individual HTC staff working at the visited sites. **290,888 (91%)** of these tests were documented in the reviewed logbooks and an additional **30,333 (9%)** could be attributed to individual providers from staff codes in the HTC registers. **268** (11%) of 2,234 providers with documented activity had tested 300 or more clients

this quarter. A dedicated full-time HTC provider is expected serve 300 clients per quarter (average of 5 clients on 60 working days per quarter). The 268 HTC providers who met or exceeded this target produced **134,272 (42%)** of the total number of tests accounted for by individual staff this quarter.

4.2.2 ART/PMTCT

Integrated HIV program supervision has included a staffing census for ART clinics since Q3 2014. This census is implemented during the site visits, indicating all staff members who actually worked at the ART clinic on the most recent clinic day. The census is designed to provide an accurate snapshot of the actual staffing of ART services each quarter. The numbers collected may be slightly lower than longer term averages, because around 80 service delivery staff are themselves participating in the supervision exercise and will not be counted as having worked in their ART clinic during the supervision period. The table below shows that total staffing levels have been fairly consistent over the last 3 quarters.

In April 2015, **600** clinicians (physicians, clinical or medical officers); **812** nurses and **838** auxiliary staff (health surveillance assistants, clerks, etc.) were working in ART clinics in Malawi.

Quarter	Clinicians	Nurses	Pharmacy Staff	Auxiliary Staff
2014 Q3	595	829	11	833
2014 Q4	616	854	11	848
2015 Q1	600	812	14	838

An estimated 2.6 million ART patient visits are currently managed at the 714 ART sites per annum, based on approximately 550,000 patients alive on ART and an average dispensing interval of 2.5 months. With 260 working days per year, an average of 10,200 patient visits are therefore managed by the ART sites per working day. At current staffing levels, this translates into an average of **17** ART patient visits per clinician and **12** per nurse per day. This approximate HRH capacity assessment does not take account of site-specific differences in patient burden and staffing levels and there are several medium and high burden sites with sub-optimal staffing. However, the national treatment program is fully decentralized to the health centre level and the program continues to devolve the growing patient burden to peripheral facilities. Since 2011, the steepest increase in ART patient numbers has been recorded at the 300 small peripheral sites that have the largest collective staffing capacity.

5 HIV Testing and Counselling Program Outputs

HTC protocols have been revised in 2013 and a new HTC register was implemented in the course of a national re-training campaign for all HTC providers between May and November 2013. Protocol revisions include:

- Clear recommendations for re-testing based on the client's test result and risk assessment
- Proper documentation of confirmatory testing for clients with a prior positive result (usually performed at enrolment into care).

This is the second HTC report based on the 2013 HTC register. The full national HTC data are presented in the **Appendix**.

5.1 HTC data

500,086 people[3] were tested and counselled for HIV between January and March 2015. **470,997 (94 %)** of these tests were performed at health facilities, **16,293 (3%)** were done in stand-alone HTC sites and **12,796 (3%)** were done outside of facilities / in the community. Out of a total of **31,484** people newly diagnosed

[3] Reports from the HTC register are based on client encounters. It is not possible to de-duplicate people who access HTC multiple times in the reporting period. However, very few individuals come for repeat testing in less than 3 months and the number of HTC encounters in one quarter is therefore assumed to represent individuals.

with HIV this quarter, **29,836 (95%)** were tested at health facilities, **1,371(4%)** at stand-alone HTC sites and **277 (1%)** in a community-based testing.

5.2 HTC access type

287,187 (57%) of people tested were patients receiving provider-initiated testing and counselling (PITC); **211,197 (42%)** accessed voluntary counselling and testing, door-to-door, community-based testing, etc. **1,702 (<1 %)** came for testing with a *Family HTC Referral Slip* (FRS) that was issued to a family member at a prior HTC encounter. Based on a total of 14,383 FRS issued to index clients this quarter, the successful referral rate for family members was **12%** (1,702/ 14,385). This is lower than in previous quarter (16%). Referral slip issuance and utilization has remained low.

5.3 Age and sex distribution among HTC clients

Out of **500,086** people tested and counselled, **33%** were males and **67 %** were females. **50%** of females were pregnant. The proportion of males (50%) and non-pregnant females (50%) was very similar, implying gender balanced access to HTC services. Pregnant women have to be excluded from the comparison of male and female access to HTC because testing among pregnant women is almost entirely provider-initiated and there is no comparable access route targeting males.

49% of all people tested and counselled were 25 years and above, **41 %** were between 15-24 years and **10%** were children below 15 years. **115,450 (23%)** accessed HTC with their partners (as a couple).

5.4 First time, repeat and confirmatory test results

The 2011 and 2014 Malawi Clinical HIV Guidelines stipulate: *All patients need a confirmatory HIV test to rule out any possibility of mix-up of test results or fraudulent access to ART: either at enrolment into pre-ART follow-up, or before starting ART if the test to confirm was not done in pre-ART. Children under 12 months starting ART with a positive DNA-PCR do not need another confirmatory test before starting ART, but all need a confirmatory rapid antibody test at age 12 and 24 months.*

157,032 (31%) accessed HTC for the first time and **343,279 (69%)** were repeat testers. Based on the cumulative number of people who accessed HTC for the first time, a total of **5,722,700** people have been tested since introduction of the *first time HTC access* indicator in July 2007.

31,491 (6%) out of all clients received a positive result for the first time. Positive rapid test results among infants (**926**) and inconclusive test results (**1385)** both accounted for **<1 %** of new results given to clients.

330,051 (96 %) of 324,054 repeat testers reported a *last negative* result. **10,565 (3%)** were reported as *previous positives* and all of these should have been classified as receiving a confirmatory test. For most of the **10,565** previous *positives*, testing was probably initiated by a health worker before enrolment into care. *Confirmatory test results* accounted for **9,983 (94%)** of *previous positive* clients. The remainder (582) may have been misclassified as *new positive* or *new inconclusive* because they were among clients who independently sought confirmation of their positive status. **9,405 (94%)** of 9,983 confirmatory tests were concordant positive and **578 (6%)** were classified as *confirmatory inconclusive*. This category includes parallel concordant negative and discordant test outcomes (Determine HIV1/2 and Uni-Gold HIV1/2 are used in parallel for confirmatory testing). This relatively high proportion of clients who did not have a concordant positive confirmation may be explained by selective confirmatory testing among clients with doubts about their previous positive status, but it underscores the importance of both routine confirmatory testing before ART initiation as well as the need to continue strengthening HTC quality assurance processes.

Figure 1: Confirmatory HIV testing coverage at ART sites in the 5 zones

Num.: total confirmatory HIV tests documented in HTC registers. Denom.: total new patients initiating ART at the site



The 9,983 confirmatory test results documented this quarter indicate that only **40%** of the 25,264 patients initiating ART this quarter received confirmatory testing and **Figure 1** shows that confirmatory testing coverage was low in all 5 zones. Only **78 (11%)** of facilities throughout the country had performed confirmatory testing for ≥75% of patients newly initiated on ART. Implementation of the confirmatory testing policy will be further reinforced over the next quarters.

6 DNA-PCR testing for Early Infant Diagnosis of HIV (EID)

DNA-PCR testing is performed at 7 labs (Mzuzu Central Hospital, Mzimba District Hospital, Kamuzu Central Hospital, Queen Elizabeth Central Hospital, DREAM Blantyre, Zomba Central Hospital and Partners in Hope, Lilongwe). EID counsellors collect infant blood samples as dried blood spots on filter paper. Health facilities are requested to maintain a standard EID DNA-PCR logbook to document EID samples and to track results. The logbook includes the dates of collection, dispatch, receipt of result from the lab and communication of the result to the mother. Supervision teams were asked to collect basic data from these logbooks.

506 (79%) of 640 sites with HIV exposed children in follow-up had collected and recorded at least 1 DNA-PCR sample during Q1 2015. A total of **8,257** DNA-PCR samples were collected and recorded. By the time the logbooks were reviewed (between 2 and 4 weeks after the end of the quarter), results had been received at the sites for **4,487 (54%)** of these specimens and **2,272 (51%)** of these results had been communicated to the mother. The proportion of results received at the sites was **71%**, **67%** and **28%** for samples collected in January, February and March, respectively. A total of **247 (6%)** results received at the sites were positive.

The **7 laboratories** dispatched DNA-PCR test results for **7,526** children in Q1 2015. This is 1,001 (12%) fewer than the number of samples recorded in the DNA-PCR logbooks at health facilities during this quarter. This discrepancy is likely due to the difference in sample collection and result dispatch dates. **6,261 (83%)** of the dispatched results were from samples collected in Q1 2015, while 1,265 (17%) were from samples collected

in the previous quarters. The median time between sample collection and dispatch of the result was **19 days**; 75% of results were dispatched between 14 and 29 days after sample collection.

3,395 (46%) of all results were from infants under 2 months old at the time of sample collection. 3,197 (43%) were 2-5 months, 731 (10%) were 6-11 months and 80 (1%) were 12 months or older when the sample was collected (date of birth was missing for 123).

Age at sample collection	Tot. Results	Positives	
<2 months	3,395	50	*1.5%*
2-5 months	3,197	97	*3.0%*
6-11 months	731	75	*10.2%*
12 months +	80	8	*10.0%*

233 (3.1%) of all results dispatched were positive. The age-specific number (%) of positive results is shown on the left. Receipt of the DNA-PCR result at the health facility is a prerequisite to updating of patient records and for appropriate clinical management. Considering the delays between sample collection and dispatch of the test result from the lab, the child's age at the time of dispatch of the result from the lab is a useful indicator for the process of early infant diagnosis and early initiation of ART for confirmed infected infants. The table below shows the distribution of ages when results were dispatched from the lab.

Age when result disp. from lab	Tot. Results	*(Col %)*	Positives	*(Col %)*
<2 months	833	*11%*	10	*4%*
2-5 months	5,538	*74%*	122	*52%*
6-11 months	874	*11%*	81	*35%*
12 months +	158	*2%*	17	*7%*
(missing date)	123	*2%*	4	*1%*
Total	7,526	***100%***	233	***100%***

Out of 233 positive results dispatched, only 10 (4%) were sent before the child was 2 months old. A total of 132 (57%) positive results were sent before the child was 6 months old and 213 (91%) were sent before the child was 12 months old. A total of 116 infants were started on ART in WHO stage 1 or 2 on the basis of confirmed HIV infection (see ART section below). This is equivalent to **54%** of the positive DNA-PCR results dispatched for children <12 months this quarter.

7 Blood Safety

The Malawi Blood Transfusion Service (MBTS) is striving to provide blood products for the entire country using voluntary non-remunerated donors and quality assured screening for transfusion transmissible infections (TTIs). For the last years, MBTS has not been able to meet the national demand and several hospitals continue to supplement or rely entirely on blood units collected from replacement donors. Complete reports from MBTS have been available throughout, but blood safety reports from health facilities have not been consistently available and it has been challenging to compile national reports relying on the data passively submitted by the sites. Therefore, the PMTCT/ART supervision teams were tasked with active collection of blood donor and cross-matching data from all visited health facilities. Some of the visited laboratories were not using the standard MOH registers and the aggregation of data for reporting may have been affected by incomplete documentation at some sites.

A total of **19,436** blood units were collected in Malawi during Q1 2015. MBTS collected **12,640 (65%)** of these, **100%** of which were screened comprehensively for the relevant TTIs (HIV, Hepatitis B, Hepatitis C, syphilis, malaria). In addition, **54** hospitals in Malawi collected a total of **6,796** units from replacement donors. **6,125 (91%)** of these units were screened for at least the 3 key TTIs (HIV, HepB and syphilis) and **2,127 (35%)** of these were also screened for HepC and malaria. This means that a total of **18,765 (97%)** of all 19,436 units collected by MBTS and from replacement donors this quarter were screened at least for HIV, HepB and syphilis. Based on the blood donor registers at the sites that collected blood from replacement donors, 12 units were screened for HIV and HepB only and 12 were screened only for HIV. 647 were screened with any other combination of tests for TTIs.

A total of **10,541** potential replacement donors were documented in the blood donor registers at the facilities and 6,796 (64%) of these ended up donating. Facilities may have used different screening algorithms

and potential donors may have been excluded on the basis of different criteria, including TTIs, blood group, haemoglobin concentration and/or clinical conditions. Testing for less prevalent TTs may have only been carried out for donors who passed the screening for more common conditions. In total, 80% of potential donors were tested for HIV, 80% for HepB, 80% for syphilis, 62% for malaria and 49% for HepC. Detailed data on individual test outcomes among all potential blood donors are presented in the Appendix.

8 Post Exposure Prophylaxis (PEP)

A total of **1,271** persons received PEP during Q1 2015. This is 10% decrease from the previous quarter (1,409).

9 Provider-Initiated Family Planning (PIFP)

The Integrated Clinical HIV Guidelines encourage health workers to routinely provide condoms to all adults in pre-ART and ART clinics. Women should also be offered at least the standard injectable contraceptive (Depo-Provera) during any pre-ART or ART visit. This policy aims to address the significant unmet need for family planning that had been observed among HIV patients in Malawi and to reduce the number of unwanted pregnancies among HIV-infected women (***PMTCT Prong 2***). HIV program reporting on PIFP is limited to women who received an injection of Depo-Provera in pre-ART and ART clinics during the last quarter. The report does not account for family planning need nor does it include women who accessed family planning services outside of HIV clinics.

Table 3: Number and % of women retained in HIV care * who were on injectable contraceptives (Depo) by the end of 2015 Q1.

	Pre-ART		ART		Both patient groups	
Zone	Tot. women	On Depo	Tot. women	On Depo	Tot. women	On Depo
NZ	682	**89** *13%*	30,926	**5,149** *17%*	31,608	**5,238** *17%*
CEZ	487	**43** *9%*	25,652	**2,136** *8%*	26,139	**2,179** *8%*
CWZ	2,889	**551** *19%*	64,701	**14,764** *23%*	67,590	**15,315** *23%*
SEZ	2,957	**774** *26%*	99,041	**23,198** *23%*	101,998	**23,972** *24%*
SWZ	4,754	**1,170** *25%*	105,198	**30,713** *29%*	109,951	**31,883** *29%*
Malawi	11,769	**2,627** *22%*	325,517	**75,960** *23%*	337,286	**78,587** *23%*

* estimated from the total number of patients retained in pre-ART and ART, multiplied by the proportions of females and adults registered

Table 3 shows that **78,587 (23%)** of 337,286 women in care received Depo-Provera from HIV clinics in Q1 2015. The SW Zone had achieved the highest coverage among women in pre-ART and ART. Patient coverage and stock availability had further improved this quarter and 582 (82%) of ART/PMTCT sites had stocks of Depo-Provera in April 2015.[4] This was mainly due to inclusion of Depo-Provera in the quarterly distribution of ARVs and other HIV commodities.

10 Cotrimoxazole Preventive Therapy (CPT)

All patients in HIV care are universally eligible for CPT in order to reduce the frequency and severity of several HIV-related diseases. Patients with confirmed HIV infection are provided lifelong CPT in pre-ART and ART clinics. CPT is also given to HIV exposed children until exposure to breast milk has stopped and HIV infection has been ruled out (usually around age 24 months). Fewer than 5% of patients are expected to require stopping of CPT due to toxicity.

[4] Many Mission hospitals do not provide family planning.

Table 4 shows that **624,739 (92%)** of 681,189 all patients in care were on CPT at the end of Q1 2015.

Table 4: Number and % of patients retained in HIV care who were on cotrimoxazole and isoniazid preventive therapy (CPT, IPT) by the end of 2015 Q1.

	CPT								IPT	
	Exp. child		Pre-ART		ART		All patient groups		Pre-ART	
Zone	Tot. pat.	On CPT	Tot. pat.	On CPT	Tot. pat.	On CPT	Tot. pat.	On CPT	Tot. pat.	On IPT
NZ	8,046	**6,570** *82%*	2,625	**2,559** *97%*	54,648	**53,075** *97%*	65,319	**62,204** *95%*	2,625	**2,190** *83%*
CEZ	7,727	**6,207** *80%*	2,065	**2,004** *97%*	44,659	**43,088** *96%*	54,451	**51,299** *94%*	2,065	**1,549** *75%*
CWZ	16,376	**14,152** *86%*	8,990	**7,182** *80%*	111,991	**95,508** *85%*	137,357	**116,841** *85%*	8,990	**6,405** *71%*
SEZ	30,576	**27,381** *90%*	11,190	**11,027** *99%*	159,317	**144,855** *91%*	201,083	**183,262** *91%*	11,190	**10,017** *90%*
SWZ	27,768	**24,057** *87%*	15,482	**14,974** *97%*	179,729	**172,102** *96%*	222,979	**211,132** *95%*	15,482	**13,972** *90%*
Malawi	90,493	**78,366** *87%*	40,352	**37,746** *94%*	550,344	**508,627** *92%*	681,189	**624,739** *92%*	40,352	**34,134** *85%*

11 TB / HIV Interventions

11.1 Intensified Case Finding (ICF)

TB is one of the most important HIV-related diseases in Malawi and a considerable proportion of (mainly early) deaths on ART are attributed to undiagnosed TB. ICF is carried out using a standard symptom checklist at every HIV patient visit. ICF outcomes are documented on HIV exposed child, pre-ART and ART patient cards, but routine M&E reporting is currently limited to ART patients in order to reduce the burden of reporting secondary cohort outcomes. It is assumed that implementation of ICF is similar in pre-ART and exposed child follow-up.

538,510 (98%) of all patients retained on ART were screened for TB at their last visit before end of March 2015. As of that visit, **5,534 (1%)** patients were new TB suspects and had presumably been referred for examination by a clinician and for TB investigations. **1,161 (<1%)** patients had confirmed TB (clinical or lab based). Out of these, **978 (84%)** were confirmed to be on TB treatment and **183 (16%)** had not yet started or had interrupted TB treatment. An excerpt from the data in the **Annex** (*Cumulative ART outcomes*) is shown below.

Current TB status among ART patients (ICF)

ICF not done (Current TB status unknown/ not circ)		**11,834**	**2%**
ICF done		**538,510**	**98%**
	TB not suspected	**531,815**	99%
	TB suspected	**5,534**	1%
	TB confirmed	**1,161**	0%
		TB confirmed, not on treatment	**183**	16%
		TB confirmed, on TB treatment	**978**	84%

11.2 Isoniazid Preventive Therapy (IPT)

All pre-ART patients with a negative screening outcome for TB symptoms are eligible for IPT. The first (large scale) distribution of isoniazid and pyridoxine for the HIV programs reached the sites during July 2012. **34,134 (85%)** of 40,352 patients retained in pre-ART were on IPT by the end of March 2015. Isoniazid was in stock at 641 facilities during the April 2015 supervision visit.

12 HIV-Related Diseases

Table 5 shows the number of patients treated for key HIV-related indicator diseases. **4,158** patients were started on TB treatment this quarter and HIV status was ascertained for **3,765 (91%)**. **1,954 (52%)** of these were HIV positive and **1,408 (72%)** of all HIV positives were already on ART when starting TB treatment. The Diflucan (fluconazole) donation program has received renewed attention and significant stocks of fluconazole were distributed to all sites in November 2012. In Q1 2015, **865** and **601** patients received Diflucan for acute cryptococcal meningitis and oesophageal candidiasis, respectively. **260** patients with Kaposi sarcoma were registered for ART in this quarter.

Table 5: Number new cases of key HIV-related diseases registered per quarter (KS = Kaposi Sarcoma, CM = cryptococcal meningitis, OC = oesophageal candidiasis).

	TB				KS *	CM *	OC *
	Tot. cases	HIV status asc.	HIV positive	Already on ART	Tot. cases	Tot. cases	Tot. cases
2014 Q2	4,258	3,773 *89%*	2,130 *56%*	1,619 *76%*	312	408	774
2014 Q3	4,692	4,368 *93%*	2,149 *49%*	1,460 *68%*	310	368	640
2014 Q4	4,204	3,989 *95%*	2,103 *53%*	1,546 *74%*	336	475	663
2015 Q1	4,158	3,765 *91%*	1,954 *52%*	1,408 *72%*	260	865	601

13 HIV-Exposed Child Follow-Up

13.1 Methods and Definition of Indicators

There are multiple entry points into HIV exposed child follow up: children of HIV infected mothers may be enrolled at birth at maternity / postnatal ward; they may be found at Under 1 or Under 5 Clinics through active screening for HIV exposure; they may be identified when presenting sick to OPD; or they may be seen with their mothers in ART follow-up. Although the targeted enrolment age is below 2 months, children may theoretically be enrolled up to 23 months of age (when HIV infection can be ruled out by rapid antibody test and breast milk exposure is likely to have stopped).

Initial registration data and details for every visit are recorded on an *Exposed Child Patient Card* and a subset of the registration data is copied in the *HIV Care Clinic (HCC) register* (one record per patient). Registration data are reported from the HCC register on a quarterly basis. Follow-up outcomes are reported monthly, selecting children who were **2, 12 and 24 months** old in the respective reporting month. Outcomes are determined from the latest visit details recorded on each card. HIV infection status is evaluated as ***known negative*** if a negative DNA-PCR or rapid test result was available at the last visit; HIV infection status is evaluated as ***known positive*** if a positive DNA-PCR result was available at any age or a positive rapid antibody test was available from age 12 months; HIV infection status is counted as ***unknown*** if HIV infection has not been confirmed and/or a negative test result pre-dated the last visit (assuming on-going HIV exposure through breast milk). All children under 24 months with confirmed HIV infection and those under 12 months with confirmed HIV infection through DNA-PCR or HIV antibody and symptoms of *presumed severe HIV disease* are ***eligible for ART***.

The main outcome indicator for the HIV exposed child follow-up program is ***HIV-free survival at 24 months of age***. This is defined as the proportion of children who were discharged as confirmed HIV uninfected by the age of 24 months.

13.2 HIV Exposed Child Registration Data

10,452 HIV exposed children were newly enrolled into follow-up during Q1 2015; **6,750 (65%)** of these were under the age of 2 months. This represents timely enrolment for **79%** of the 8,499 known HIV exposed children discharged from maternity this quarter. The total number of new enrolments (10,452) exceeds by 1,953 (23%) the total number of known HIV exposed children discharged from maternity (8,499). This apparent discrepancy may be explained by delayed enrolment of infants born in previous quarters; by double-counting of infants who transferred between sites; or by identification and enrolment of additional HIV exposed infants after birth. Overall, enrolment into follow-up for known HIV exposed infants appears to be almost complete.

The documentation of follow-up outcomes, particularly the updating of DNA-PCR results on patient cards, remained incomplete at several sites. This has led to an underreporting of ascertainment of HIV status among the 2 month old cohort.

13.3 Birth Cohort Outcomes

There were **8,342** infants in the **2 month age cohort**. **3,116 (37%)** had received a DNA-PCR result. **56 (2%)** of these were confirmed HIV infected. An additional **24** infants were diagnosed with *presumed severe HIV disease,* which means that a total of **80** infants were eligible for ART. **48 (60%)** of these had started ART. The proportion of positives starting ART has decreased from the previous quarter (80%). Out of the entire 2-month age cohort, **7,580 (92%)** were retained in exposed child follow-up, **48 (1%)** had started ART and **30 (<1%)** were discharged confirmed uninfected[5]. **13(<1%)** were known to have died and **526 (6%)** had been lost to follow-up.

There were **9,230** children in the **12 month age cohort**. Current HIV infection status was known for **4,082 (44%)** children (DNA-PCR or rapid antibody test) and **187 (5%)** of these were confirmed HIV infected. **20 (<1%)** additional children had been diagnosed with *presumed severe HIV disease,* which means that a total of **207** children were eligible for ART. **172 (83%)** had started ART. Out of the entire age cohort, **6,746 (75%)** were retained in exposed child follow-up, **172 (2%)** had started ART and **100 (<1%)** were discharged confirmed uninfected.[5] **1,911 (21%)** were lost to follow-up and **76 (<1%)** were known to have died.

There were **8,999** children in the **24 month age cohort**. Current HIV infection status was known for **4,102 (46%)** children (DNA-PCR or rapid antibody test) and **242 (6%)** of these were confirmed HIV infected. **7** additional children had been diagnosed with *presumed severe HIV disease,* which means that a total of **249** children were eligible for ART. **242 (97%)** of these had started ART. Out of the entire age cohort, **1,007 (12%)** were retained in exposed child follow-up, **242 (3%)** had started ART and **3,757 (43%)** were discharged confirmed uninfected. **3,596 (41%)** were lost to follow-up and **111 (1%)** were known to have died.

Confirmed HIV-free survival at age 24 months in this quarter remained implausibly low at **43%.** This was related to the fact that only 46% in this cohort had a known HIV status. 4,897 (54%) children were classified as '*current HIV infection status unknown'* and many of these may be among the 3,596 children lost to follow-up and the 111 children who had died. However, 1,007 (12%) were retained in follow-up beyond age 24 months and a final rapid test was not available for these children, possibly due to continued breast feeding. There are still problems with scheduled HIV testing (and documentation of test results) at 6 weeks, 12 and 24 months of age.

[5] A small number of children may be rightfully discharged as 'confirmed uninfected' by 2 or 12 months of age, provided that HIV exposure through breast milk has definitely stopped (e.g. maternal death) and a negative HIV test was obtained at least 6 weeks thereafter.

14 Pre-ART

14.1 Pre-ART Registration Data

A total of **6,223** patients were newly registered for pre-ART follow-up in Q1 2015. **461 (4%)** of these were children aged 5-14 years. The number of new pre-ART enrolments continued to decline from the previous quarter (6,483 total, 485 children) due to the introduction of relaxed ART eligibility criteria in April 2014. Several sites had already established pre-ART services before July 2011 and the cumulative number of pre-ART patients ever registered was **182,555**.

14.2 Cumulative Pre-ART Follow-up Outcomes

40,352 (22%) of all patients ever registered were retained in pre-ART follow-up by the end of March 2015; **94,850 (51%)** had started ART; **44,778 (25%)** had been lost to follow-up; **2,011 (1%)** were known to have died. The proportion of patients starting ART is bound to increase in the cumulative pre-ART cohort analysis over time. Based on a subtraction of cumulative outcomes from the previous quarter, **5,150** pre-ART patients started ART during Q1 2015; **1,642** were lost to follow-up and **262** died.

CPT coverage among pre-ART patients was **94%** in Q1 2015 while IPT coverage remained at **85%**. **2,627 (22%)** of 11,769 women had received Depo-Provera from their pre-ART clinic. Further details on CPT, Depo-Provera and IPT are presented in **Tables 3** and **4** in the sections above.

15 PMTCT / ART

The implementation of ***PMTCT Option B+*** has effectively integrated PMTCT and ART services. The program aims to initiate lifelong ART for all HIV infected women as early as possible in pregnancy. ART may be started and continued at ANC, labour and delivery, and at ART clinics. All infants born to HIV-infected women are supposed to start daily nevirapine prophylaxis for the first 6 weeks of life. Nevirapine syrup is given to women at ANC at the earliest opportunity to take home with instructions how to give it to the new-born.

15.1 Data Sources and Reporting Methods

New standard M&E tools for ANC and maternity were implemented in January 2010 and revised in Q2 2012 to reflect the Option B+ policy. ANC and maternity clinic registers and reporting forms include patient management information and all relevant data elements for the maternal and child health and HIV programs. The ANC register was specifically designed to avoid data duplication that previously affected PMTCT reports from ANC due to the inability to account for individual women's outcomes in the course of multiple visits. The cohort reporting system is designed to aggregate women's outcome data after they have completed their ANC visits. Data from ANC and maternity are collated and presented separately because records do not allow identification of individual women and hence are subject to double counting if not separated.

All patients starting ART are recorded using standard program monitoring tools (ART patient treatment cards and ART clinic registers). **ART baseline data** for all patients registered are reported each quarter from ART clinic registers. **ART outcomes** of all patients ever registered are reported after reviewing the cards of all new patients and of those who were on ART at the end of the previous quarter, updating the status of patients who have subsequently died, stopped or been lost to follow-up. Secondary outcomes such as current regimen, CPT status, side effects, adherence and TB status are reported for all patients retained on ART.

ART scale-up has resulted in a growing proportion of HIV-infected women who are already on ART when getting pregnant. Implementation of *Option B+* will further increase ART coverage in this group. **Maternal ART coverage** is estimated from the number of pregnant women who were already on ART when getting pregnant (**maternity reports**) ***plus*** those who newly started ART when pregnant (**ART reports**).

Maternity reports capture ART status at the time of delivery (up to the time of discharge from the postnatal ward). The timing of ART initiation is categorized into: (any time) before pregnancy; during 1st / 2nd trimester; during 3rd trimester; during labour. About 97% of pregnant women in Malawi attend ANC, but only 83% of women in the general population deliver at a health facility in Malawi. Maternity reports therefore have the potential for undercounting the number of mothers and infants receiving ARVs. However, there is evidence from ANC and maternity reports that almost all of the known HIV infected women deliver at health facilities. ARV coverage among known positives is therefore reliably calculated from maternity reports. Women admitted at maternity who are referred to another facility before / after delivery are double-counted in aggregated maternity data. Assuming the probability of referral is independent of ART status, the number of women already on ART when getting pregnant is therefore **adjusted** by the overall proportion of referrals among women admitted to maternity.

ART program reports capture pregnancy (and breastfeeding) status at the time of *ART initiation*, providing information on the number of new women starting ART while pregnant (or while breastfeeding). ART reports do not capture women who become pregnant after starting ART. For the estimation of maternal ART coverage, the number of women starting ART in pregnancy is **adjusted for:**

a) Double-counting of women starting ART in pregnancy and subsequently transferring to another site. These women are counted multiple times as 'pregnant at the time of starting ART' in the quarterly ART cohort reports because the disaggregation of age, sex and reason for starting ART applies to all patients newly registered in the quarter, including transfers in. Separate *ART 'survival' analyses* are collected each quarter for women started under Option B+. The proportion of women transferred within 12 months of registration is used to adjust the quarterly number of pregnant women starting ART for transfers.

b) *Failed ART initiation* is thought to be the main underlying reason for early loss to follow-up among the Option B+ cohort. Patients are recorded on patient cards and in clinic registers when the first supply of ARVs is dispensed and all new entrants are counted as ART initiations in the quarterly ART cohort report. Recent operational studies indicate that most pregnant women lost to follow-up within the first 6 months never return after this first dispensing visit and many of these may have never actually started taking ART. The proportion of women lost to follow-up in the 6-month survival analysis is therefore used to adjust the number of pregnant women starting ART in the quarterly ART cohort reports for *failed initiations.*

Infant PMTCT coverage is estimated from maternity reports, based on the number of infants born to known HIV-infected women and discharged alive who started nevirapine prophylaxis.

Coverage is calculated by dividing the number of patients served by population denominators. The denominators are derived from expected pregnancies based on population projections and HIV prevalence from epidemiological surveillance (source: 2014 Spectrum model for Malawi). There are an estimated 13,317 HIV infected pregnant women in the population per quarter (1/4 of 53,268 in 2014).[6]

15.2 ARV Coverage among Pregnant / Breastfeeding Women and Exposed Infants

11,583 (80%) of the estimated 13,317 HIV infected pregnant women in Malawi this quarter were on ART. This is based on **6,266** [7] women at maternity who were already on ART when getting pregnant and **4,317** [8] women who newly initiated ART in pregnancy.

[6] 2014 Spectrum estimates based on 2014 definition of eligibility for ART in Malawi (CD4<500, Option B+, UT for U5).

[7] 6,596 women who started ART before pregnancy admitted at maternity; reduced by 5% to adjust for double-counting of 6,602 referrals among 129,368 total admissions.

[8] 6,078 women registered at ART clinics who were pregnant at the time of starting ART; a) 8% are discounted to adjust for double-counting of transfers based on 664 of 8,392 women who transferred within 12 months of registration (12 month Option B+ survival analysis); b) 22.8% are discounted to account for presumed failed ART initiations based on 1,835 of 8,036 women lost to follow-up within 6 months of registration (6 month Option B+ survival analysis).

An additional **1,584** [9] breastfeeding women started ART due to ***Option B+*** (in WHO clinical stage 1 or 2), bringing the total number newly started on ART under *Option B+* to **5,901**. Most women starting ART while breastfeeding were probably identified late in maternity or early in the postnatal period, but this group may also include some women who re-initiated after interrupting ART in pregnancy. **7,866** infants were confirmed to have started NVP prophylaxis at maternity.

Figure 2 shows the transition from prophylactic ARV regimens for HIV infected mothers to universal ART under ***Option B+*** (registration data; not adjusted as above). The (less effective) single dose NVP regimen and AZT combination prophylaxis had been phased out by April 2012. The average number of pregnant women registered for ART each quarter **increased almost 6-fold** from **1,221** in the 12 month period before introduction of Option B+ to an average of **7,000** since Q4 2011.

Figure 2: Transition from prophylactic ARV regimens for PMTCT to Option B+ in Malawi

Women who moved to Option B+ from sdNVP / AZT were double counted between Q3 2011 - Q1 2012. It is likely that <12,000 total women were on ARVs during these quarters. Data on women already on ART when getting pregnant are only available from Q2 2012.



15.3 HIV Services at ANC

The full national data from ANC are presented in the **Appendix**.

158,328 women attended ANC for their first visit between January and March 2015. This is 93% of the estimated 170,732 pregnant women in the 2015 population during one quarter.[10]

[9] 1,722 women registered at ART clinics who were breastfeeding at the time of starting ART; reduced by 8% to adjust for double-counting of transfers based on 664 out of 8,392 women who transferred within 12 months of registration (12 month Option B+ survival analysis). Failed ART initiations are thought to be less common among this group, so no further adjustment is made.

[10] Estimated as ¼ of 682,926 births projected for 2014.
Malawi National Statistics Office. (2008). Malawi Population Projections 2008-2030.
Retrieved from http://www.nsomalawi.mw/images/stories/data_on_line/demography/census_2008/MainReport/ThematicReports/Population Projections Malawi.pdf

The following report covers the outcomes of the **164,275** women who started ANC between July and September 2014 and who had finished ANC between January and March 2015. **13,245 (8%)** of these started ANC in their first trimester. **8,958 (5%)** were tested for syphilis at ANC and **891 (10%)** were syphilis positive. The low testing rate probably explains the higher (10%) than expected proportion (<1%) of positives as the testing was likely selective of those suspected to be positive. Only **35,500 (22%)** of women in this cohort attended the minimum of 4 focussed ANC visits.

15.3.1 HIV Ascertainment at ANC

148,838 (91%) of ANC attendees had their HIV status ascertained. This is a slight improvement to the previous quarter (89%). Out of all women whose HIV status was ascertained, **12,274 (8%)** presented with a valid documented previous HIV test result and **136,564 (92%)** received a new HIV test result at ANC. A total of **11,727 (7.8%)** women were found HIV positive. This is lower than the estimated 11% HIV prevalence among pregnant women in the 2010 ANC sentinel surveillance survey but consistent with the latest Spectrum projections (7.8% HIV prevalence among pregnant women in 2014).[6]

15.3.2 ARV Coverage at ANC

10,821 (92%) of (known) HIV infected women attending ANC received ART. This represents **81%** coverage of the estimated 13,317 HIV positive pregnant women per quarter at the population level.

Of the **10,821** ANC women who were known to receive ART, **5,403 (50%)** were already on ART when starting ANC, **4,238 (39%)** initiated before 28 weeks of pregnancy and **1,180 (11%)** initiated during the last trimester of pregnancy. **10,828 (92%)** of HIV infected women at ANC were on Cotrimoxazole Preventive Therapy.

9,454 (81%) of known HIV infected women attending ANC received the infant dose of ARVs (nevirapine syrup) to take home.

15.4 HIV Services at Maternity

The full national data from maternity are presented in the **Appendix**.

Between January and March 2015, **122,766** women were admitted for delivery to maternity; **6,602** of these were referred to another facility before delivery, resulting in **129,368** total admissions to maternity during Q1 2015. Out of all admissions, **114,970 (95%)** delivered at health facilities, while **6,183 (5%)** had already delivered before reaching a facility. The **114,970** facility deliveries represent **67%** of the estimated 170,732 quarterly deliveries in the population in 2015 which is less than the 83% reported in the Integrated Household Survey Report of 2010-2011.

A total of **112,425 (94%)** deliveries were conducted by skilled birth attendants, **485 (<1%)** by paramedical staff and **6,026 (5%)** were not attended by any of the above (probably mainly among women who delivered before reaching maternity). **15,121 (12%)** of women developed obstetric complications. The most common leading complications were obstructed / prolonged labour (**4,840** cases) and post-partum haemorrhage (**1,670** cases). A total of **121,153** babies were born, **117,192 (97%)** were singletons and **3,961 (3%)** were twins/multiples. There were **119,103 (98%)** live births and **2,050 (2%)** stillbirths. **118,003 (99%)** of babies born alive were discharged alive and **1,100 (1%)** died before discharge. **118,944 (>99%)** of women were discharged alive and **92 (<1%)** women died before discharge, which is equivalent to a maternal mortality ratio of **76 per 100,000** live births among women attending maternity.

15.4.1 HIV Ascertainment at Maternity

120,651 (96%) women had their HIV status ascertained at maternity. Out of these, **118,081 (98%)** presented with a valid previous HIV test result and **2,570 (2%)** received a new HIV test result. A total of **9,142 (8%)** women were HIV positive and **111,509 (92%)** were negative. The **120,651** women whose HIV status was ascertained at maternity represent **71%** of the expected 170,732 women delivering in the population.

HIV exposure status was ascertained for **113,891 (97%)** out of 118,003 babies born and discharged alive. **8,499 (7%)** of these were born to a known HIV positive mother.

15.4.2 ARV Coverage at Maternity

A total of **8,860 (97%)** of known HIV infected women admitted to maternity received ART. Out of these, **6,596 (74%)** had started ART before pregnancy, **1,164 (13%)** initiated ART during the 1st or 2nd trimester, **902 (10%)** initiated during the 3rd trimester and **198 (2%)** initiated ART at maternity.

A total of **7,866 (93%)** of 8,499 infants who were known HIV exposed and discharged alive started daily NVP prophylaxis at maternity. This represents **59%** coverage of the estimated 13,317 HIV exposed infants born in the population in this quarter.

16 ART Access and Follow-Up Outcomes

The full national data from the ART Program are shown in the **Appendix**.

16.1 New ART Registrations during Q1 2015

By the end of March 2015, there were **714 static ART sites** in Malawi, managed by government, mission, NGOs and the private sector. Out of these, **95** were ART facilities in the private sector, charging a nominal MK500 per monthly prescription of drugs per patient.

The electronic reporting system supported by MSF in Chiradzulu district experienced a prolonged technical fault and accurate ART cohort reports for the 10 affected facilities could not be obtained in time for this report. It was therefore decided to use the previous quarter's results for these 10 sites. This has led to some inconsistencies in the data presented.

Implementation of the Malawi Integrated Clinical HIV Guidelines started in July 2011, triggering a massive surge in new ART initiations (see **Figure 3**). **25,270** patients initiated ART in Q1 2015 and **6,280** patients were registered as a transfer in (already on treatment; 20% out of all 32,033 clinic registrations). These numbers are similar to previous quarter.

Among all new registrations **37%** were males, **63%** were females and **6,122 (30%)** of females were pregnant. For some of the **6,078** who had started under ***Option B+*** and who transferred to sites with electronic medical records system, the status at registration was updated to 'no longer pregnant', leading to an apparent discrepancy with the number of women recorded as pregnant. An additional **1,722** women in WHO stage 1 or 2 were started because of breastfeeding, bringing the total number of women registered as started under ***Option B+*** [11] to **7,800**.

[11] Universal ART for all HIV infected pregnant and breastfeeding women in WHO stage 1 or 2, independent of CD4 count

Figure 3: Patients newly inititated on ART and total ART clinic registrations per quarter

Total ART clinic registrations include patients who transferred between sites. This results in double counting of patients at the national level. For 'patients newly initiated on ART' every patient is only counted once.



A total of **20,189 (63%)** of all patients registered started in WHO stage 1 or 2 and **11,456 (57%)** of these started due a low CD4 count. **9,768 (31%)** of patients registered started in WHO stage 3 and **1,653 (5%)** started in stage 4.

2,883 children were registered at ART sites in Q1 2015. **708** of these were registered under the expanded policy of universal ART for children aged 12-59 months in WHO stage 1 or 2, independent of CD4 count. **109** children started ART with presumed severe HIV disease, which was lower than the previous quarter (136). **116** infants in WHO stage 1 or 2 who started due to confirmed HIV infection through DNA-PCR, which is slightly higher than the previous quarter (106). Early paediatric ART access has remained below expectations, but the relatively low number is consistent with reduced transmission rates due to Option B+: considering that 8,499 HIV exposed infants were identified at maternity and assuming a 2% transmission rate among the 97% of HIV positive mothers at maternity who received ART (and 20% transmission in the 3% who did not receive ART)[12], only about 216 of these known HIV exposed infants may have been infected perinatally during Q1 2015. However, considering the projected 1,560 new infant HIV infections in the 2015 population per quarter[6], early infant treatment coverage remains low at an estimated **7%** (116 / 1,560). The most significant bottleneck for early infant treatment remains the identification of HIV infected pregnant / breastfeeding women.

1,033 (3%) out of all ART clinic registrations were patients with TB: **592 (2%)** had a current and **441 (1%)** a recent history of TB. **260 (1%)** of patients registered had Kaposi's sarcoma.

16.2 Cumulative ART Registrations up March 2015

By the end of March 2015, there were a cumulative total of **994,006** clinic registrations, representing **795,144 (80%)** patients who newly initiated ART and **187,893 (19%)** patients who transferred between clinics. **10,969**

[12] UNAIDS Reference Group on Estimates Modelling and Projections (2011). Working paper on mother-to-child-transmission rates for use in Spectrum. Geneva, UNAIDS.

(1%) out of all clinic registrations were patients who re-initiated ART after treatment interruption. Out of all registrations, **36**% were males and **64**% were females, **91**% were adults and **9**% were children (<15 years). Private sector clinics accounted for **30,116** (3.0%) of total patient registrations.

16.3 ART Outcomes

551,566 patients were alive on ART by the end of March 2014. This number includes 1,232 patients assumed to be in transit between sites (50% of the 2,464 new transfers out this quarter).

Out of the **795,144** patients ever initiated on ART, **551,566 (69%)** were retained alive on ART, **75,536 (9%)** were known to have died, **178,071 (22%)** were lost to follow-up and **3,109** (<1%) were known to have stopped ART. An estimated **503,604** adults and **47,962** children (<15 years) were alive on ART by the end of March 2015**.**

Figure 4 Patients alive on ART at the end of each quarter, stratified by size of facility (number of patients alive on ART)



Figure 4 shows the increase of patients alive on ART by the end of each quarter. The number of patients alive on ART **increased by 16,370** in Q1 of 2015. **Figure 4** also illustrates the ongoing decentralization of Malawi's ART program. From Q3 2011, the greatest increase in ART patient numbers was seen at sites with fewer than 500 patients alive on ART. By the end of March 2015, **49%** of the national ART patient cohort was in care at sites with fewer than 2,000 patients.

Figure 5: Quarterly rates of ART drop out (ART stop, defaulters and deaths)

Numerator: new ART stops, new defaulters and new deaths in the respective quarter

Denominator: total patients retained alive at the end of the previous quarter plus new patients registered in the respective quarter)



Figure 5 shows the considerable decrease of ART drop-out rates since the start of the national program. There were **2,102** new deaths, **10,402** new defaulters in Q1 2015. This translates into a quarterly death rate of **0.4%** and a defaulter rate of **1.9%** among the patients alive and on treatment in this quarter. These rates were similar to the previous quarter. Based on previous operational studies, about half of the patients in stage 3 and 4 who are classified as lost to follow-up are thought to have died. There is also an indication that 10-15% of pregnant women who were registered as 'initiated on ART' under Option B+ may have never actually started taking ARVs due to inadequate preparation at ANC. Importantly, the ascertainment of loss to follow-up requires updating of patient treatment cards after analysis of the most recent dispensing visit. Any lack in rigour in this process will lead to a misclassification of patients who have been lost to follow-up as 'retained alive on ART'. The cumulative total number of patients known to have stopped declined from the previous quarter, probably due to missing data from flood-affected sites.

By end of March 2015, a cumulative **75,536 (9%)** patients were known to have died **178,071 (22%)** were lost to follow-up and **3109** (**<1%**) were known to have **stopped ART.**

Figure 6: Patients starting ART in WHO stage 4 and deaths in the first 3 months after ART initiation. (Shown as proportions among new patients registered each quarter)



Figure 6 shows the considerable decline in **early mortality** since the start of the program. In Q2 of 2006 11% of new patients died within the first 3 months after ART initiation. There has been a remarkable decline in early mortality and the lowest point thus far has been reached in Q4 2013. The decrease in early mortality is probably mainly due to earlier ART initiation (patients in WHO stage 2 with a CD4 count below the threshold or in stage 3). It correlates well with the decline in the proportion of patients starting ART in WHO clinical stage 4 from 25% in 2005 Q2 to **5%** in Q4 2014. Slight fluctuations in the calculated early mortality rates are mainly due to inconsistent classification of month of death at the sites with electronic patient record systems. The 2014 guidelines are expected to further reduce early mortality, as more patients will be started in WHO stage 1 and 2 (universal ART for HIV infected pregnant and breastfeeding women and children under 5 years).

16.4 ART Cohort Survival Analysis

A 12, 24, 36, 48, 60, 72, 84, 96 and 108-month **'cohort outcome survival analysis'** was conducted for patients registered in Q1 of 2006, 2007, 2008, 2009, 2010, 2011, 2012, 2013 and 2014, respectively. A separate 12-month cohort outcome analysis was conducted for children who were under 15 years at the time of ART initiation and who registered for ART in Q1 2014. For the 12th time, a further subgroup analysis was done for women who started ART under ***Option B+*** during Q1 2013, Q4 2014 and Q3 2014. **76% of adults** and **77% of children** were retained alive on ART after 12 months on treatment. This is similar to the previous quarter and remains below the WHO target of 85%.

The majority of patients classified as lost to follow-up are likely to have stopped/ interrupted ART, but others will have transferred to another facility without notifying the previous site. Actual retention rates are thought to be about **10%** higher due to misclassification of 'silent transfers' as 'defaulters' in clinic-based survival/retention analysis. A population-based study in Karonga district with individual linkage showed that

92% of patients started in 2011-2012 were retained after 12 months on ART while routine monitoring data showed **79%** retention rates for the same period.[13]

Figure 7 shows the continuous improvement of long-term treatment outcomes over time. **58%** and **52%** of patients registered 5 and 7 years ago had been retained alive on ART.

Figure 7: Group cohort survival analysis: Proportion of patients retained alive on ART 12, 24, 36, 48, 60, 72, 84 and 96 months after ART initiation



6-month group cohort survival outcomes were known for **8,508** (99%) of the 8,782 women registered as having started ART under *Option B+* in Q3 2014. [14] This number represents 472 (6%) women who transferred out and are therefore double counted and **8,036 (94%)** patients not transferred. **6,148 (77%)** of these were retained at 6 months after registration. **1,835 (97%)** of those not retained were lost to follow-up, **12 (1%)** were known to have stopped ART and **41 (2%)** were known to have died.

12-month group cohort survival outcomes were known for **8,392 (97%)** of the 8,707 women registered as having started ART under *Option B+* in Q1 2014. [14] This number represents **664 (8%)** women who transferred out and are therefore double counted and **7,728 (92%)** patients not transferred. **5,601 (72%)** of these were retained at 12 months after registration. **2,029 (95%)** of those not retained were lost to follow-up, **32 (2%)** were known to have stopped ART and **66 (5%)** were known to have died.

24-month group cohort survival outcomes were known for **8,714 (95%)** out of the 9,096 women registered as having started ART under *Option B+* in Q1 2013. [14] This number represents **863 (10%)** women who transferred out and are therefore double counted and **7,851 (90%)** patients not transferred. **5,411 (69%)** of these were retained at 24 months after registration. **2,314 (95%)** of those not retained were lost to follow-up, **42 (2%)** were known to have stopped ART and **84 (3%)** were known to have died.

[13] Koole, O., Houben, R. M. G. J., Mzembe, T., Van Boeckel, T. P., Kayange, M., Jahn, A., Crampin, A. C. (2014). Improved retention of patients starting antiretroviral treatment in Karonga District, northern Malawi, 2005-2012. Journal of Acquired Immune Deficiency Syndromes (2014), 67(1), e27–33. doi:10.1097/QAI.0000000000000252

[14] Group cohort survival analyses were not available from some sites with electronic data systems. 'Reason for starting' may be reclassified for some patients, leading to minor inconsistencies in patients included in group cohort survival analyses.

2,289 (25%) of the women in the 24 month Option B+ survival cohort had initiated ART in the breastfeeding period and **2,110 (23%)** started in the third trimester / in labour; considering the 24 month median breastfeeding period in Malawi (2010 MDHS), more than half of the women in this cohort can be assumed to have stopped breastfeeding. The **69% and 70% retention rate at 24 and 36 months** after ART initiation confirms that a high proportion of women started under Option B+ **remain on ART beyond the cessation of breastfeeding**.

The 6-month retention rate is the same as in the previous quarter. These are satisfactory results. Most of the women lost to follow-up failed to return after their first visit and many of these may have never actually started ART due to inadequate counselling and preparation in the initial phase of implementation. The program is examining the different modes of delivery closely to further improve uptake and retention on *Option B+*.

6 month survival OptionB+

Survival and retention in ART program *

ART cohort registration group outcomes

Total ART clinic registrations	8,508	100%
Transfers out (double counted)	472	6%
Total not transferred out (patients in cohort)	8,036	94%
Total alive on ART	6,148	77%
Total not retained	1,888	23%
Defaulted	1,835	97%
Stopped ART	12	1%
Died	41	2%

12 month survival OptionB+

Survival and retention in ART program *

ART cohort registration group outcomes

Total ART clinic registrations	8,392	100%
Transfers out (double counted)	664	8%
Total not transferred out (patients in cohort)	7,728	92%
Total alive on ART	5,601	72%
Total not retained	2,127	28%
Defaulted	2,029	95%
Stopped ART	32	2%
Died	66	3%

24 month survival OptionB+

Survival and retention in ART program *

ART cohort registration group outcomes

Total ART clinic registrations	8,714	100%
Transfers out (double counted)	863	10%
Total not transferred out (patients in cohort)	7,851	90%
Total alive on ART	5,411	69%
Total not retained	2,440	31%
Defaulted	2,314	95%
Stopped ART	42	2%
Died	84	3%

36 month survival OptionB+

Survival and retention in ART program *

ART cohort registration group outcomes

Total ART clinic registrations	11,951	100%
Transfers out (double counted)	1,466	12%
Total not transferred out (patients in cohort)	10,485	88%
Total alive on ART	7,315	70%
Total not retained	3,170	30%
Defaulted	2,916	92%
Stopped ART	53	2%
Died	201	6%

16.4.1 Secondary outcomes of patients retained on ART

Secondary outcomes are known for the **550,344** patients alive on ART who remained at their sites at end of the quarter.

ART Regimens

543,630 (99%) of patients were on first line and **6,309 (1%)** were on second line regimens; **405 (<1%)** were on non-standard regimens. Non-standard regimens are not necessarily substandard regimens and include patients continuing an ART regimen that was started outside Malawi, patients in research programmes and patients in specialist care.

Among patients on first line regimens, **26,627 (5%)** were on paediatric formulations and **25,158 (94%)** of these were on the new standard first line for children (regimen 2P: AZT/3TC/NVP). By the end of March 2015, **478,644 (93%)** of patients on adult first line were receiving regimen **5A** (tenofovir / lamivudine / efavirenz). **26,840 (5%)** were on regimen 2A (zidovudine / lamivudine / nevirapine), which was the main alternative regimen for patients with stavudine side-effects before transition to regimen 5A and **1,206 (<1%)** were on regimen 1A (stavudine / lamivudine / nevirapine).

Adherence to ART

Pill counts and the number of missed doses were documented for **545,532 (99%)** out of all patients retained on ART and **493,486 (90%)** of these were classified as >95% adherent in Q1 2015. Manual estimation of adherence from pill counts is practically difficult and classification can be misleading. The ART program has switched to a direct evaluation of doses missed in 2010 to improve on accuracy of adherence assessment and plausible adherence levels are recorded with this method. However, there have also been persistent challenges with the analysis of adherence levels at sites with Electronic Data Systems (EDS) and adherence data from several of these sites could not be included in this report.

ART Side Effects

528,700 (96%) patients on ART had information on drug side effects documented at their last clinic visit before end of March 2015. **3,955 (1%)** of these had side-effects. The prevalence of side effects seems to have stabilized at very low levels following the full transition to regimen 5A (tenofovir / lamivudine / efavirenz) that started in July 2013.

16.5 Viral Load (VL) Monitoring

The National Treatment Program has started rolling out routine VL monitoring for patients on ART to facilitate early detection of treatment failure and timely switching to second line ART. Routine VL monitoring is scheduled at 6 months after ART initiation, at 2 years and every 24 months thereafter. Additional targeted VL testing may be carried out for patients with clinically suspected treatment failure. During Q1 2015, **8** laboratories in the national program provided VL testing for patients enrolled at the 8 respective facilities and associated sites. A total of **19,062** VL results were produced at these labs between January and March 2015. The number of VL results per lab was: Thyolo District Hospital: 3,954; Partners in Hope (Lilongwe): 3,390; Kamuzu CH (Lilongwe): 3,678; DREAM (Blantyre): 2,875; Zomba CH (Zomba): 1,261; Mzuzu CH (Mzimba): 1,623; QECH (Blantyre): 1,165; Mzimba DH (Mzimba): 1,116.

Reason	0-999		1000-4999		*5000+*		*Total*
Routine	**15,555**	*86%*	**644**	*4%*	**1,808**	*10%*	**18,007**
Targeted	**343**	*54%*	**43**	*7%*	**244**	*39%*	**630**
Other/unk	**219**	*51%*	**39**	*9%*	**168**	*39%*	**426**
Total	**16,116**	*85%*	**726**	*4%*	**2,220**	*12%*	**19,062**

18,007 (94%) of all VL samples were classified as *routine scheduled.* This is equivalent to **26%** of the estimated 70,000 ART patients passing a VL monitoring milestone this quarter. **630 (3%)** of samples were classified as *targeted (suspected treatment failure / repeat)* and for **426 (2%)** the reason for the sample was other or not specified. **15,974 (85%)** of all results were undetectable

/ below 1,000 copies/ml. The proportion of results with 5,000+ copies was higher among targeted samples (39%) than among *routine* samples (10%). VL monitoring outputs are expected to increase significantly over the next quarters.

17 TB / HIV Management

Approximately **80%** of HIV infected TB patients were receiving ART in Q1 2015. This estimate is based on the following triangulation of TB and ART program data:

TB Program Data: A total of **4,158** TB patients were registered during Q1 2015. Assuming an average HIV prevalence of 60% among TB patients, **2,495** TB patients were HIV positive and therefore in need of ART. Given that **1,408** TB patients registered were already on ART at the time of starting TB treatment, 2,495 – 1,408 = **1,087** TB patients needed to initiate ART.

ART Program Data: An estimated **592** patients[15] started ART with a current or recent episode of TB in Q1 2015. This is **54%** (592 of 1087) of the TB patients who needed to start ART. This means that a total of 1,408 + 592 = **2,000 (80%)** of the estimated 2,495 HIV infected TB patients were receiving ART in Q1 2014.

TB program report *

TB clinic registrations

Total TB patients registered	4,158	100%

HIV status ascertainment

HIV status not ascertained	393	9%
HIV status ascertained	3,765	91%
HIV negative	1,811	48%
HIV positive	1,954	52%
Already on ART	1,408	72%
Not on ART when starting TB treatment	546	28%

TB / ART program triangulation *

HIV-burden among TB patients (estimated)

HIV negative (est. 40%)	1,663	40%
HIV positive (est. 60%) in need of ART	2,495	60%
Not on ART	274	11%
Total on ART (coverage)	2,221	89%
Already on ART (TB prog)	1,408	63%
Started ART within 24m of TB diagnosis (ART prog)	813	37%
ART initiations with current TB (ART prog)	466	57%
ART initiations after recent TB (ART prog)	347	43%

18 STI Treatment

STI reports were actively collected during the Integrated HIV Program Supervision exercise for the 8th time this quarter. This decision was taken due to the persistent challenges with 'passive' reporting based on data aggregated by the district STI coordinators. This quarter, supervision teams collected STI data from 697 out of 928 facilities offering STI management according to the *2013-14 Service Provision Assessment*[16] in Malawi. The site-level reports included here may therefore only represent 73% of all STI services in Malawi. The supervision teams re-emphasized the importance of complete and accurate documentation at the sites and

[15] 21% of the 1,012 ART patients who were registered this quarter with a recent or current episode of TB at the time of ART initiation were assumed to be transfers and were subtracted to adjust for double-counting.

[16] Ministry of Health, & ICF International. (2015). Malawi Service Provision Assessment (SPA) 2013-14. Lilongwe, Malawi and Rockville, Maryland, USA. Retrieved from http://dhsprogram.com/pubs/pdf/SPA20/SPA20.pdf

the data quality is expected to improve with resumption of regular site supervision for the STI program. The complete set of STI program data collected is included in the Appendix.

18.1 Access to STI treatment and coverage

Based on the data collected at the facilities, a total of **54,897** STI cases were treated in Q1 2015. Considering the 73% site-level completeness of reporting, this number is estimated to represent a total of **75,201** STI cases treated. This is equivalent to **76% STI treatment coverage** of the expected 98,600 STI cases in the population.

Out of **54,897** documented clients treated, **21,850** (40%) were male and **33,047** (60%) were female. **4,238** (13%) of female STI clients were pregnant. **36,823** clients (67%) were 25 years and above, **13,176** (24%) were 20-24 years and **4,898** (9%) were under 20 years old.

18.2 Client Type and STI History

48,409 (88%) of clients were symptomatic and **6,488** (12%) were asymptomatic (treated as partners). Among symptomatic clients, **44,127** (91%) of were index cases and **4,282** (9%) were partners. A total of **19,731** partner notification slips were issued, equivalent to an average of 0.45 slips per index case. Considering the 19,731 partner notification slips issued, **55%** (10,770) of those notified presented to the clinic. **40,691** (74%) of clients presented with their first lifetime episode of STI, **9,608** (18%) clients reported to have had an STI more than 3 months ago and **4,598** (8%) of clients reported having had an STI within the last three months. Re-occurrence of an STI after a recent episode may be due to re-infection or treatment failure.

18.3 HIV Status

HIV status was ascertained for **26,317** (48%) clients and **7,587** (29%) of these were HIV positive. **1,607** (21%) of positives were identified through a new test initiated at the STI clinic, while **5,980** (79%) presented with a documented previous positive HIV test result. **4,622** (77%) of clients with a previous positive HIV test result were on ART.

The rate of HIV status ascertainment at STI clinics remained low. This is likely due to poor implementation of provider initiated testing and counselling, combined with weak back-referral systems which may lead to incomplete documentation of new HIV test results at the STI clinics. It is worth noting that a substantial proportion of clients who are aware of their HIV infection present with a new episode of an STI. This may suggest poor translation of positive living strategies promoted during counselling, but could also be due to the increased risk of recurrence of HSV-2 and balanitis among HIV-infected clients.

18.4 STI Syndromes

The most common syndrome was abnormal vaginal discharge (AVD) with **17,974** (30%) cases, followed by urethral discharge (UD, **13,008** cases), genital ulcers (GUD, **10,229** cases) and lower abdominal pain (LAP, **9,246** cases). Similar to previous reports, balanitis, bubo, warts and neonatal conjunctivitis each accounted for 1 – 2% of cases.

18.5 Referrals

Given the high risk of recent HIV infection among STI clients, all clients with unknown status and those with a new negative test result should be referred for (repeat) HIV testing and counselling. **17,766 (37%)** of the 47,580 STI clients with unknown or new negative test result were referred for repeat HTC. **856 (53%)** of 1,607 clients who were newly tested HIV positive were referred for ART eligibility assessment.

19 Supply of HIV Program Commodities

19.1 Quantification and procurement planning

The program conducted the quarterly quantification and procurement planning for all HIV commodities in January 2015 prior to submission of the Global Fund Concept Note in January 2015. This formed the basis for the HIV Commodity Quantification assumptions for ARVs, OI medicines and selected laboratory commodities for the period ending December 2017.

During Q1 2015, there was a marked improvement noted with the procurement and delivery of 5A (tenofovir / lamivudine / efavirenz) resulting into 5.1 and 5.2 months of stock in the warehouse and health facility levels respectively as per the April 2015 National Stock status report. **478,644 (93%)** of 517,003 patients receiving first line adult formulation ART were on this preferred regimen by the end of March 2015.

During Q1 2015, ARVs and medicines for opportunistic infections worth $13.2 million were received by the Bollore Africa Logistics managed warehouses dedicated for Department of HIV and National Malaria Control Program commodities. This comprised of Tenofovir/Lamivudine/Efavirenz 300/300/600mg (Regimen 5A; 83% of the value of adult ARVs) and other ARV formulations (17% of the value for all medicines received during the period).

To maintain adequate stocks in the pipeline and hence ensure uninterrupted supply for subsequent orders, the Ministry has continued processing HIV commodity orders for ARVs, OI, RDTs and other related commodities through Partnership for Supply Chain Management (ARVs and RDTs) and IDA Foundation (laboratory commodities and medicines for opportunistic infections).

19.2 Quarterly distribution of HIV Commodities

The scheduled quarterly distribution of HIV commodities (Distribution Round 21) took place between mid-February to mid-March. 43 HIV and Antimalarial commodities (Antimalarials, ARVs, OI, STI medicines and laboratory commodities) were distributed to 726 health facilities. Both Malaria RDTs and HIV test kits were also distributed to individual health facilities to ensure adequate stocks and uninterrupted testing services at all sites. This was the second successful consolidated distribution for HIV and anti-malarial commodities.

During Q1 2015, the logistics team at the Department of HIV and AIDS also coordinated a total of over 1,500 individual commodity transactions between ART sites. This is to avert stock outs and/or expiry. All these transactions are managed using the HIV Department Supply Chain Hot Line, a toll free facility that was set up to facilitate communication between the health facilities and the central level. Health workers are able to communicate supply chain and other HIV commodities related issues that need to be resolved by the technical team at the department in a timely manner.

19.3 Quarterly logistics support during Q1 2015 ART/PMTCT supervision

The logistics section supported development of the distribution list for HIV commodities for 726 ART/PMTCT Sites for Distribution Rounds 22 and 23 to cover the consumption periods of Q2 (April-June 2015) and Q3 (July-Sep 2015).

HTSS and HIV seconded Supply Chain and Logistics Officers also provided stock management support at 292 sites during the Q1 2015 integrated ART/PMTCT Site visits supervision. This is aimed at ensuring that physical inventory is conducted at all sites. Mentoring in stock management was provided at health facilities with poor storage practices. There is a marked improvement in the logistics management of ARVs and medicines for opportunistic infections. Health care providers have continued to use RDT daily activity registers and relocation books for registration of redistributed commodities to health facilities for which authorization codes must be obtained as a commodity tracking measure. This enables the program foster accountability before commodities are issued from an ART/PMTCT and HTC sites.

19.4 National Stock Status of HIV Commodities

Physical stock counts for ARVs and other medicines for HIV-related diseases were performed at all sites during the supervision visits in April 2015. **Table 6** shows the total medicine stocks found at the sites and the estimated consumption periods. Health facility stocks of the key adult and paediatric regimens were estimated to last until August 2015.

Minimum stocks of TDF/3TC 300/300mg and AZT/3TC 300/150mg are maintained at all sites for post-exposure prophylaxis (PEP) and the total stocks at the sites therefore far exceeds the actual consumption from patients using this regimen in alternative ART regimens or as PEP.

Commodities in high volumes include stavudine and efavirenz containing regimens at all levels mainly driven by the transition to tenofovir based regimens. The program continues to monitor the trend of patients on stavudine and efavirenz containing regimens to inform future procurements for alternative regimens such as Abacavir/Lamivudine formulations given that Stavudine containing regimens will be phased out in the long run.

478,644 patients were on regimen 5A, which was 12,764 **(2.6%)** less than projected in the previous forecast for the end of this quarter **(**491,408**).** The national ART program forecast and quantification was updated in July 2015 to inform procurement planning and budgeting for HIV commodities for the period ending December 2017.

Table 6: Total stocks of HIV program commodities at all sites visited during the 2015 Q1 supportive site supervision. Stock positions are from the date of the visit (between 1-4 weeks after the end of the quarter). Warehouse stock positions are from 20/04/2015

Inventory unit	Item	Sites with any Stock	Total Physical Stock		Consump-tion/ Month	Months of Stock *	
			At Sites	In Warehouse		At Sites	Wareh.
tins	ABC / 3TC 60 / 30mg tins (60 tabs)	148	13,336	22,337	3,939	3.4	5.7
	ABC / 3TC 600 / 300mg tins (30 tabs)	10	92	5,000	563	0.2	8.9
	ATV / r 300 / 100mg tins (30 tabs)	165	9,568	4,581	5,486	1.7	0.8
	AZT / 3TC / NVP 300 / 150 / 200mg tins (60 tabs)	619	115,306	328,034	26,881	4.3	12.2
	AZT / 3TC / NVP 60 / 30 / 50mg tins (60 tabs)	633	328,396	476,554	63,778	5.1	7.5
	AZT / 3TC 300 / 150mg tins (60 tabs)	485	12,267	28,421	2,299	5.3	12.4
	AZT / 3TC 60 / 30mg tins (60 tabs)	572	20,741	24,138	2,621	7.9	9.2
	d4T / 3TC / NVP 30 / 150 / 200mg tins (60 tabs)	120	16,658	11,752	1,208	13.8	9.7
	d4T / 3TC 30 / 150mg tins (60 tabs)	217	12,758	764	124	102.9	6.2
	d4T / 3TC 6 / 30mg tins (60 tabs)	202	2,409				
	EFV 200mg tins (90 tabs)	146	2,245	3,525	363	6.2	9.7
	EFV 600mg tins (30 tabs)	209	25,180	5,777	824	30.6	7.0
	LPV / r 100 / 25mg tins (60 tabs)	66	9,417	2,223	2,421	3.9	0.9
	LPV / r 200 / 50mg tins (120 tabs)	59	580	51	431	1.3	0.1
	NVP 200mg tins (60 tabs)	369	27,014	20,988	6,667	4.1	3.1
	NVP 50mg tins (60 tabs)	136	11,627	1,800	1,265	9.2	1.4
	TDF / 3TC / EFV 300 / 300 / 600mg tins (30 tabs)	705	2,348,035	2,292,120	481,021	4.9	4.8
	TDF / 3TC 300 / 300mg tins (30 tabs)	600	54,743	139,981	11,408	4.8	12.3
bottles	Fluconazole (Diflucan) 50mg / 5ml bottles (35 ml)	9	847	2,000	90	9.4	22.1
	Gentian violet 25g bottles (1 each)	423	8,696		1,188	7.3	
	NVP 10mg/ml bottles (25 ml)	593	123,301	88,441	9,252	13.3	9.6
vials	Benzathine Penicillin 1.44g vials (50 each)	600	427,086		38,254	11.2	
	Bleomycine 15,000IU vials (1 each)	24	29,939				
	Ceftriaxone 1g vials (50 each)	428	151,650		103,254	1.5	
	Depo-Provera 150mg/1ml vials (25 each)	581	873,004	964,775	262,720	3.3	3.7
	Gentamicin 80mg / 2ml vials (50 each)	462	266,730		97,166	2.7	
	Vincristine 1mg / 1ml vials (1 each)	79	40,333	31,193	3,120	12.9	10.0
tabs	Aciclovir 200mg blist packs (500 tabs)	578	4,083,679	3,119,150	622,412	6.6	5.0
	Amitriptylline 25mg tins (500 tabs)	232	731,975		118,590	6.2	
	Azithromycin 500mg blist packs (3 tabs)	282	77,255		10,272	7.5	
	Ciprofloxacin 500mg blist packs (100 tabs)	431	2,033,571		294,410	6.9	
	Clotrimazole 500mg boxes (1 each)	339	798,545		37,844	21.1	
	Codeine 30mg tins (100 tabs)	146	608,175		48,726	12.5	
	Cotrimoxazole 100 / 20mg blist packs (1000 tabs)	616	34,354,897	31,233,000	6,514,064	5.3	4.8
	Cotrimoxazole 400 / 80mg tins (1000 tabs)	630	40,549,951	8,679,000	16,322,639	2.5	0.5
	Cotrimoxazole 960mg blist packs (1000 tabs)	546	19,998,763	21,140,000	17,356,290	1.2	1.2
	Doxycycline 100mg tins (1000 tabs)	463	8,776,661	587,000	4,362,250	2.0	0.1
	Erythromycin 250mg tins (1000 tabs)	320	3,367,863	1,498,000	3,902,474	0.9	0.4
	Fluconazole (Diflucan) 200mg tins (28 tabs)	328	391,195	714,280	63,925	6.1	11.2
	Fluconazole (generic) 200mg tins (100 tabs)	33	60,168				
	Ibuprofen 200mg tins (100 tabs)	219	1,310,948		834,358	1.6	
	Isoniazid (H) 100mg blist packs (100 tabs)	138	148,262		147,127	1.0	
	Isoniazid (H) 300mg tins (1000 tabs)	641	11,622,028	14,108,000	1,076,838	10.8	13.1
	Metronidazole 200mg tins (1000 tabs)	415	8,049,973		4,738,819	1.7	
	Morphine 10mg blist packs (60 tabs)	134	1,007,298		212,625	4.7	
	Pyridoxine 25mg tins (100 tabs)	143	1,909,017		1,149,434	1.7	
	Pyridoxine 50mg tins (1000 tabs)	536	9,582,564	9,147,000	1,149,434	8.3	8.0
sheets	ART pat. card adult (yellow) bundles (100 sheets	602	169,002	198,700	9,713	17.4	20.5
	ART pat. card paed. (blue) bundles (100 sheets)	592	87,270		961	90.8	
	Exposed child card (pink) bundles (50 sheets)	588	62,797		3,483	18.0	
	Polythene sleeve bundles (100 sheets)	566	202,011	156,000	16,230	12.4	9.6
	Pre-ART pat. card (green) bundles (100 sheets)	592	141,204		2,073	68.1	
tests	DBS kit (filter paper, lancet, etc.) bundles (20 eac	528	51,742		31,681	1.6	
	Determine HIV1/2 boxes (100 each)	646	626,338	1,245,900	106,658	5.9	11.7
	Determine syphilis boxes (100 each)	341	93,696				
	Uni-Gold HIV1/2 boxes (20 each)	467	39,119	68,440	13,855	2.8	4.9
pieces	Condoms female boxes (1000 each)	379	967,166		182,716	5.3	
	Condoms male boxes (144 each)	591	13,499,310	6,315,552	4,372,760	3.1	1.4

* 'Consumption per month' and 'Months of stock' for ARVs, CPT, INH and HIV test kits are based on the respective patient-regimen groups in the standard service reports. Estimates are based on the number of patients on the respective regimen at the end of the quarter evaluated and do not account for potential (positive or negative) growth. Facility stock positions for OI and STI drugs include HIV Program and other supply sources. Total national consumption and MoS estimates are used for these commodity groups. 'Months of stock' is calculated from the day of the physical stock count, which is on average 1 month after the end of the quarter.

20 Training and Mentoring

ART/PMTCT trainings

487 nurses and **124** clinicians were refreshed in the 2014 national guidelines in the quarter. **22** tutors were also refreshed. Cumulatively, **5,483** health workers (providers) have been refreshed.

255 were newly trained in the 2014 national clinical guidelines. 187 were nurse and 68 clinicians.

STI training

124 clinicians and nurses have been trained in STI syndromic management approach between January and March, 2015.

HIV Testing and Counselling trainings

19 participants were trained and certified in initial HTC training during this quarter.

21 Participants in Q1 2015 Supervision (Site visits 13 April-1 May 2015)

Absalom Kaunda (CO, MOH, Mzimba DHO)
Alefa Fikira (CMT, MOH)
Amos Makwaya (CO, MOH)
Andraida Mtoseni (Nurse, MOH)
Andrew Dimba (, NTP)
Angela Nkhoma (Nurse, MOH)
Annie Biza (Nurse, MDF)
Batoni Upindi (TB Zonal Supervisor, MOH)
Benson Isake (, NTP)
Bonface Chione (CO, Lighthouse)
Cecilia Sambakunsi (Logistics Fellow, HIV Dept)
Chifundo Makuluni (Nurse, MOH)
Chikayiko Majamanda (Nurse, MOH)
Chikondi Harrison (, Logistics)
Chimwemwe Francis Mkandawire (IT Fellow, Dept for HIV and AIDS)
Chisuwo Laphiod (, moh)
Chrissy Lizengo (, MOH)
Christopher Mkwezalamba (CO, MOH)
Cornelius Kang'ombe (, NTP)
Dalitso Midiani (PMTCT Officer, MOH)
Davie Maseko (CO, SOS)
Dorica Chirwa (Logistic officer, MOH)
Edith Thaulo (Nurse, MOH)
Eliza Mahimanya (Logistics Officer, MOH)
Emmanuel Nkonde (, NTP)
Erik Mittochi (CO (ART coord), MOH)
Evans Kagwira (TB Zonal Supervisor, MOH)
Everista Mkandawire (Nurse, MOH)
Fainala Muyila (Nurse, MOH)
Freser Nkawa (, moh)
Geoffrey Makhalira (, NTP)
Gerald Zomba (Program Officer, Dept for HIV and AIDS)
Grant Gondwe (, NTP)
Hannock Matupi (ARV clinician, MOH, Rumphi DH)
Harrison Tembo (CO, MOH)
Henry Kanyerere (TB/HIV Program Officer, MOH)
James Mwambene (CO, Diginitas)
Jane Chiwoko (Nurse, Lighthouse)
Jean Kayamba (Nurse, MOH)
Jesse Lobeni (Nurse, MOH)
Joe Gumulira (CO, MOH)
John Kabichi (CO, MOH)
Judith Ntopa (Nurse, Cobbe Barracks)
Juliana Soko (ARV nurse, MOH, Livingstonia MH)
Justice Kaphiri (, NTP)
Kelvin Makina (Logistics, Kasungu)
Knox Banda (TB Zonal Supervisor, MOH)
Kuzani Mbendera (, NTP)
Lameck Mlauzi (, NTP(MOH))
Lameck Mzava (, NTP)
Levi Mugala (, MOH)
Limbani Kadzuwa (Nurse, MOH)
Lincy Chalunda (CO, MOH)
Lindiwe Zaina (Logistics, HIV Dept)
Lucy Banda (, MOH)
Macleod Piringu (ART CORDINATOR, MOH)
Margaret Katumbi (Nurse, MOH)
Martin Katanga (CO, MOH)
Mary Gosten (MA, MOH)
Mary Kamiza (TB Zonal Supervisor, NTP)
Mathilda Kamanga (Nurse, Army)
Maxon Phiri (, MOH)
Mercy Makaika (Nurse, MOH)
Michael Eliya (PMTCT Program Officer, MOH)
Mirriam Thindwa (Clinician, Limbe H/C)
Monica Simfukwe (Nurse, MOH, Chintheche RH)
Nelson Nanchinga (, MOH)
Noel Mphasa (TB Zonal Supervisor, NTP)
Offrey Mnduwira (CO, Police)
Oscar Kasiyamphanje (Nurse, CHAM)
Patrick Paul J M Chirwa (TB Zonal Supervisor, NTP)
Paul Nyasulu (CO, Dept of HIV & AIDS)
Pax Mkupani (Logistics Fellow, MOH)
Peter Chimphero (CO, MOH)
Peter Donda (CO, Dedza DH)
Peter Mwamlima (, Public)
Relia Mandindi (, Public)
Relia Nkhata Mandindi (Logistics, HIV Dept)
Rhoda Banda (, NTP)
Rodrick Kaulere (CO, CHAM (Sister Tereza))
Rose Maviko (Nurse, Limbe HC)
Roseby Malombe (Nurse, CHAM)
Rudolf Banda (, NTP)
Ruth Deula (Nurse, CHAM)
Sabina Phiri (Nurse, MOH)
Salome Chiwewe (Nurse, MOH, Ntchisi DH)
Sidder Hambisa (ENM, MOH)
Simon Makombe (ART officer, MOH, Department of HIV and AIDS)
Stanley Ngoma (CO, MOH)
Stony Mbiriyawanda (, MOH)
Stuart Chuka (CO, MBCA)
Symon Chiumia (, MOH)
Tadzinenani Thawe (, Lighthouse)
Timothy Mwenyedini (MA, MOH)
William Mtonga (CO, CHAM)
Yusuf Bhamu (HIV Fellow, HIV Dept)
Zengani Chirwa (TA, MOH, Department of HIV and AIDS)

Report compiled by:
Frank Chimbwandira (MO/Head of Dept. for HIV and AIDS)
Austin Mnthambala (MO/Deputy of Dept. for HIV and AIDS
Simon Makombe (ART Officer, Dept. for HIV and AIDS)
Eustice Mhango (ART Officer, Dept. for HIV and AIDS)
Michael Eliya (PMTCT Officer, Dept. for HIV and AIDS)
Dalitso Midiani (PMTCT Officer, Dept. for HIV and AIDS)
Lucius Ng'omang'oma (HTC Officer, Dept. for HIV and AIDS)
Andreas Jahn (MO/TA, Dept. for HIV and AIDS)
Caroline Ntale (TA Logistics, MOH, Dept of HIV and AIDS)
Andrew Mganga (M&E Fellow, Dept. for HIV and AIDS)
Paul Nyasulu (PMTCT/ART Officer, Dept. for HIV and AIDS)
Josep Kasola (HTC Officer, Dept. for HIV and AIDS
Khumbo Ngona (HTC Officer, Dept. for HIV and AIDS)
Amos Makwaya (STI Program Officer, Dept. for HIV and AIDS)
Stone Mbiriyawanda (M&E Officer, CMED/ Dept. for HIV/AIDS)
Gerald Zomba (PMTCT/ART Officer, Dept. for HIV and AIDS)
Chimwemwe Mkandawire (IT Fellow, Dept. for HIV and AIDS)

We thank all facility staff for their sincere welcome and co-operation with the HIV Department and its partners during these supportive visits. We congratulate all staff for their excellent work.

28th August 2015

22 Appendix (Full National HIV Program Data)

Antenatal Care

Malawi (national)

2015 Q1 (1st month of quarter, 2nd month of quarter, 3rd month of quarter)

New ANC registrations in reporting period *

Women with first visit in reporting period

New women registered	**158,328**	**100%**

ANC cohort analysis *

Total women completing ANC in the reporting period

Total women in booking cohort	**164,275**	**100%**

Visits per woman

Women with 1 visit	**35,838**	**22%**
Women with 2 visits	**44,129**	**27%**
Women with 3 visits	**48,808**	**30%**
Women with 4 visits	**29,176**	**18%**
Women with 5+ visits	**6,324**	**4%**

Trimester of first visit

Started ANC 0-12 wks	**13,245**	**8%**
Started ANC 13+ wks	**151,030**	**92%**

Pre-eclampsia

No pre-eclampsia	**161,372**	**98%**
Pre-eclampsia	**2,903**	**2%**

TTV doses

0-1 TTV doses	**80,336**	**49%**
2+ TTV doses	**83,939**	**51%**

SP tablets

0 SP doses	**18,563**	**11%**
1 SP dose (1 x 3 tabs)	**44,914**	**27%**
6+ SP tablets (2 x 3 tabs)	**100,798**	**61%**

FeFo tablets

0-119 FeFo tablets	**131,984**	**80%**
120+ FeFo tablets	**32,291**	**20%**

Albendazole (Deworming)

0 Albend. doses	**30,445**	**18%**
1 Albend. dose	**134,472**	**82%**

ITN (bednets)

No ITN	**36,325**	**22%**
ITN received	**128,628**	**78%**

Syphilis status

Not tested for syphilis	**155,317**	**95%**
Tested for syphilis	**8,958**	**5%**
Syphilis negative	**8,067**	90%
Syphilis positive	**891**	10%

* Subgroups may not add to 100% due to rounding

Antenatal Care

Malawi (national)

2015 Q1 (1st month of quarter, 2nd month of quarter, 3rd month of quarter)

ANC cohort analysis

*

HIV status ascertainment

HIV status not ascertained	**15,437**	**9%**
HIV status ascertained	**148,838**	**91%**
Valid previous test result	**12,274**	8%
Previous negative	**6,329**	52%
Previous positive	**5,945**	48%
New test at ANC	**136,564**	92%
New negative	**130,782**	96%
New positive	**5,782**	4%

HIV status summary

Total women HIV negative	**137,111**	**92%**
Total women HIV positive	**11,727**	**8%**

CPT status (among HIV pos)

Not on CPT	**899**	**8%**
On CPT	**10,828**	**92%**

Final PMTCT regimen mother

No ARVs	**906**	**8%**
Any ARVs	**10,821**	**92%**
ART (by time of initiation)	**10,821**	100%
Already on ART when starting ANC	**5,403**	50%
Started ART at 0-27 weeks of pregnancy	**4,238**	39%
Started ART at 28+ weeks of preg.	**1,180**	11%

Baby's ARVs dispensed

No ARVs dispensed for infant	**2,273**	**19%**
ARVs dispensed for infant	**9,454**	**81%**

* Subgroups may not add to 100% due to rounding

ART cohort analysis

Malawi (national)

2015 Q1 (Cumulative)

Registration details

*

ART clinic registrations

Total ART clinic registrations	**994,006**	**100%**

Registration type

First time ART initiations (total patients)	**795,144**	**80%**
ART re-initiations	**10,969**	**1%**
ART transfers in	**187,893**	**19%**

Sex

Males	**357,688**	**36%**
Females	**636,318**	**64%**
Non-pregnant	**522,091**	82%
Pregnant	**114,227**	18%

Age at ART initiation

Adults 15+ yrs	**907,571**	**91%**
Children 0-14 yrs	**86,435**	**9%**
Children 2-14 yrs	**66,663**	77%
Children below 24 mths	**19,772**	23%

Reason for starting ART

Presumed severe HIV Disease	**3,420**	**0%**
Confirmed HIV infection	**990,586**	**100%**
WHO stage 1 or 2	**428,297**	43%
Total lymphocytes <threshold	**1,135**	0%
CD4 below threshold	**285,809**	67%
CD4 unknown or >threshold	**141,353**	33%
PCR infants	**2,808**	2%
Children 12-59 mths	**6,310**	4%
Pregnant women	**96,117**	68%
Breastfeeding mothers	**36,118**	26%
WHO stage 3	**455,403**	46%
WHO stage 4	**101,099**	10%
Unknown / reason outside of guidelines	**5,787**	1%

TB at ART initiation

Never TB / TB > 24 months ago	**922,989**	**93%**
TB within the last 24 months	**36,272**	**4%**
Current episode of TB	**34,745**	**3%**

Kaposi's sarcoma at ART initiation

No KS	**974,860**	**98%**
Patients with KS	**19,146**	**2%**

* Subgroups may not add to 100% due to rounding

ART cohort analysis

Malawi (national)

2015 Q1 (Cumulative)

ART outcomes

*

Primary follow-up outcomes

Total alive on ART		**549,397**	**68%**
	Alive on ART at site of last registration	**550,344**	100%
	ART patients in transit between sites	**-947**	0%
Defaulted		**178,071**	**22%**
Stopped ART		**3,109**	**0%**
Total died		**75,536**	**9%**
	Died month 1	**18,707**	25%
	Died month 2	**11,878**	16%
	Died month 3	**6,966**	9%
	Died month 4+	**37,985**	50%

Transfers between sites

Total not transferred out		**807,060**	**81%**
Transferred out		**186,946**	**19%**

ART regimens

First line regimens		**543,630**	**99%**
	Adult formulation	**517,003**	95%
		Regimen 0A	**158**	0%
		Regimen 1A	**1,206**	0%
		Regimen 2A	**26,840**	5%
		Regimen 3A	**181**	0%
		Regimen 4A	**703**	0%
		Regimen 5A	**478,644**	93%
		Regimen 6A	**9,271**	2%
	Paed. formulation	**26,627**	5%
		Regimen 0P	**391**	1%
		Regimen 1P	**159**	1%
		Regimen 2P	**25,158**	94%
		Regimen 3P	**459**	2%
		Regimen 4P	**460**	2%
Second line regimens		**6,309**	**1%**
	Adult formulation	**5,499**	87%
		Regimen 7A	**4,327**	79%
		Regimen 8A	**1,172**	21%
	Paed. Formulation	**810**	13%
		Regimen 9P	**810**	100%
Other regimen (adult / paed)		**405**	**0%**

Adherence

Adherence unknown (not recorded)		**4,812**	**1%**
Adherence recorded		**545,532**	**99%**
	0-3 doses missed	**493,486**	90%
	4+ doses missed	**52,046**	10%

ART side effects

Side effects unknown (not recorded)		**21,644**	**4%**
Side effects recorded		**528,700**	**96%**
	No side effects	**524,745**	99%
	Any side effects	**3,955**	1%

* Subgroups may not add to 100% due to rounding

ART cohort analysis

Malawi (national)

2015 Q1 (Cumulative)

ART outcomes

*

Current TB status among ART patients (ICF)

ICF not done (Current TB status unknown/ not circ)		**11,834**	**2%**
ICF done		**538,510**	**98%**
	TB not suspected	**531,815**	99%
	TB suspected	**5,534**	1%
	TB confirmed	**1,161**	0%
		TB confirmed, not on treatment	**183**	16%
		TB confirmed, on TB treatment	**978**	84%

* Subgroups may not add to 100% due to rounding

ART cohort analysis

Malawi (national)

2015 Q1 (Quarter)

Registration details

*

ART clinic registrations

Total ART clinic registrations	**32,033**	**100%**

Registration type

First time ART initiations (total patients)	**25,270**	**79%**
ART re-initiations	**483**	**2%**
ART transfers in	**6,280**	**20%**

Sex

Males	**11,735**	**37%**
Females	**20,298**	**63%**
Non-pregnant	**14,176**	70%
Pregnant	**6,122**	30%

Age at ART initiation

Adults 15+ yrs	**29,150**	**91%**
Children 0-14 yrs	**2,883**	**9%**
Children 2-14 yrs	**2,172**	75%
Children below 24 mths	**711**	25%

Reason for starting ART

Presumed severe HIV Disease	**109**	**0%**
Confirmed HIV infection	**31,924**	**100%**
WHO stage 1 or 2	**20,189**	63%
Total lymphocytes <threshold	**109**	1%
CD4 below threshold	**11,456**	57%
CD4 unknown or >threshold	**8,624**	43%
PCR infants	**116**	1%
Children 12-59 mths	**708**	8%
Pregnant women	**6,078**	70%
Breastfeeding mothers	**1,722**	20%
WHO stage 3	**9,768**	31%
WHO stage 4	**1,653**	5%
Unknown / reason outside of guidelines	**314**	1%

TB at ART initiation

Never TB / TB > 24 months ago	**31,000**	**97%**
TB within the last 24 months	**441**	**1%**
Current episode of TB	**592**	**2%**

Kaposi's sarcoma at ART initiation

No KS	**31,773**	**99%**
Patients with KS	**260**	**1%**

* Subgroups may not add to 100% due to rounding

Blood safety

Malawi (national)

2015 Q1 (1st month of quarter, 2nd month of quarter, 3rd month of quarter)

Infect. disease screening among potential donors *

HIV screening

HIV testing not done	**2,067**	**20%**
Tested for HIV	**8,474**	**80%**
HIV negative	**8,006**	94%
HIV positive	**468**	6%

Hepatitis B screening

HepB testing not done	**2,093**	**20%**
Tested for Hepatitis B	**8,448**	**80%**
HepB Negative	**8,075**	96%
HepB Positive	**373**	4%

Hepatitis C screening

HepC testing not done	**5,427**	**51%**
Tested for Hepatitis C	**5,114**	**49%**
HepC Negative	**5,053**	99%
HepC Positive	**61**	1%

Syphilis screening

Syphilis testing not done	**2,096**	**20%**
Tested for Syphilis	**8,445**	**80%**
Syphilis Negative	**8,163**	97%
Syphilis Positive	**282**	3%

Malaria screening

Malaria testing not done	**4,034**	**38%**
Tested for malaria	**6,507**	**62%**
Malaria Negative	**6,014**	92%
Malaria Positive	**493**	8%

Summary screening outcome

Not donated	**3,745**	**36%**
Donated	**6,796**	**64%**
Screened for at least HIV, HepB and syphilis	**6,125**	90%
Screened for HIV, HepB, HepC, Syphilis, Malaria	**2,127**	35%
Screened for HIV, HepB, Syphilis	**3,998**	65%
Screened for HIV, HepB	**12**	0%
Screened for HIV only	**12**	0%
Screened with any other combination of tests	**647**	10%

Cross-matching report *

Blood group typing (for units and patients)

Total blood group typing done	**22,499**	**100%**

Blood units cross-matched (by source)

Total blood units cross-matched	**17,062**	**100%**
Total units from MBTS (estimated)	**10,266**	60%
Total units from replacement donors	**6,796**	40%

Blood units cross-matched by patient group

Units cross-matched for maternity	**2,565**	**15%**
Units cross-matched for paediatrics	**7,368**	**43%**
Units cross-matched for other ward	**7,129**	**42%**

* Subgroups may not add to 100% due to rounding

Blood safety

Malawi (national)

2015 Q1 (1st month of quarter, 2nd month of quarter, 3rd month of quarter)

Cross-matching report

*

Transfusion reactions

Units transfused without adverse events	**17,048**	**100%**
Units with suspected transfusion reactions	**10**	**0%**
Units with confirmed transfusion reactions	**4**	**0%**

* Subgroups may not add to 100% due to rounding

HIV exposed child follow-up

Malawi (national)

2015 Q1 (1st month of quarter, 2nd month of quarter, 3rd month of quarter)

Age 2 months

Age cohort outcomes

*

Total children in birth cohort

Total children registered	**8,342**	**100%**

CPT status

On CPT	**7,550**	**91%**
Not on CPT	**792**	**9%**

HIV status

Current HIV infection status unknown	**5,226**	**63%**
HIV infection not confirmed, not ART eligible	**5,202**	100%
HIV infection not confirmed, ART eligible (PSHD)	**24**	0%
Current HIV infection status known	**3,116**	**37%**
Confirmed not infected	**3,060**	98%
Confirmed infected (ART eligible)	**56**	2%

ART eligibility summary

Not eligible for ART	**8,262**	**99%**
ART eligible	**80**	**1%**
ART not initiated	**32**	40%
Initiated ART	**48**	60%

Primary follow-up outcome

Discharged uninfected	**30**	**0%**
Continue follow-up	**7,580**	**92%**
Started ART	**48**	**1%**
Defaulted	**526**	**6%**
Died	**13**	**0%**

Transfers between sites

Total not transferred out	**8,197**	**98%**
Transferred out	**145**	**2%**

Age 12 months

Age cohort outcomes

*

Total children in birth cohort

Total children registered	**9,230**	**100%**

CPT status

On CPT	**6,782**	**73%**
Not on CPT	**2,448**	**27%**

HIV status

Current HIV infection status unknown	**5,148**	**56%**
HIV infection not confirmed, not ART eligible	**5,128**	100%
HIV infection not confirmed, ART eligible (PSHD)	**20**	0%
Current HIV infection status known	**4,082**	**44%**
Confirmed not infected	**3,895**	95%
Confirmed infected (ART eligible)	**187**	5%

* Subgroups may not add to 100% due to rounding

HIV exposed child follow-up

Malawi (national)

2015 Q1 (1st month of quarter, 2nd month of quarter, 3rd month of quarter)

Age cohort outcomes

*

ART eligibility summary

Not eligible for ART	**9,023**	**98%**
ART eligible	**207**	**2%**
ART not initiated	**35**	17%
Initiated ART	**172**	83%

Primary follow-up outcome

Discharged uninfected	**100**	**1%**
Continue follow-up	**6,746**	**75%**
Started ART	**172**	**2%**
Defaulted	**1,911**	**21%**
Died	**76**	**1%**

Transfers between sites

Total not transferred out	**9,005**	**98%**
Transferred out	**225**	**2%**

Age 24 months

Age cohort outcomes

*

Total children in birth cohort

Total children registered	**8,999**	**100%**

CPT status

On CPT	**1,249**	**14%**
Not on CPT	**7,750**	**86%**

HIV status

Current HIV infection status unknown	**4,897**	**54%**
HIV infection not confirmed, not ART eligible	**4,890**	100%
HIV infection not confirmed, ART eligible (PSHD)	**7**	0%
Current HIV infection status known	**4,102**	**46%**
Confirmed not infected	**3,860**	94%
Confirmed infected (ART eligible)	**242**	6%

ART eligibility summary

Not eligible for ART	**8,750**	**97%**
ART eligible	**249**	**3%**
ART not initiated	**7**	3%
Initiated ART	**242**	97%

Primary follow-up outcome

Discharged uninfected	**3,757**	**43%**
Continue follow-up	**1,007**	**12%**
Started ART	**242**	**3%**
Defaulted	**3,596**	**41%**
Died	**111**	**1%**

Transfers between sites

Total not transferred out	**8,713**	**97%**
Transferred out	**286**	**3%**

* Subgroups may not add to 100% due to rounding

HIV Care Clinic (HCC) cohort analysis

Malawi (national)

2015 Q1 (Quarter)

Registration details

*

HCC clinic registrations

Total HCC registrations	**16,675**	**100%**

Registration type

Patients enrolled first time	**15,887**	**95%**
Patients re-enrolled	**49**	**0%**
Patients transferred in	**739**	**4%**

Sex

Males (all ages)	**7,944**	**48%**
Females (all ages)	**8,731**	**52%**
Non-pregnant	**8,713**	100%
Pregnant	**18**	0%

Age at registration

Adults 15+ yrs	**5,811**	**35%**
Children 0-14 yrs	**10,864**	**65%**
Children 24 months - 14 years	**461**	4%
Children below 24 months (exposed children)	**10,403**	96%
Children 2 - below 24 months	**3,653**	35%
Infants below 2 months	**6,750**	65%

Reason for HCC registration

Exposed infants	**10,452**	**63%**
Confirmed infected patients (pre-ART)	**6,223**	**37%**

* Subgroups may not add to 100% due to rounding

HIV Care Clinic (HCC) cohort analysis

Malawi (national)

2015 Q1 (Cumulative)

Registration details

*

HCC clinic registrations

Total HCC registrations	**340,001**	**100%**

Registration type

Patients enrolled first time	**328,247**	**97%**
Patients re-enrolled	**1,082**	**0%**
Patients transferred in	**10,672**	**3%**

Sex

Males (all ages)	**146,297**	**43%**
Females (all ages)	**193,704**	**57%**
Non-pregnant	**192,706**	99%
Pregnant	**998**	1%

Age at registration

Adults 15+ yrs	**173,144**	**51%**
Children 0-14 yrs	**166,857**	**49%**
Children 24 months - 14 years	**15,888**	10%
Children below 24 months (exposed children)	**150,969**	90%
Children 2 - below 24 months	**72,750**	48%
Infants below 2 months	**78,219**	52%

Reason for HCC registration

Exposed infants	**150,704**	**44%**
Confirmed infected patients (pre-ART)	**189,297**	**56%**

Pre-ART follow-up outcome

*

Primary follow-up outcomes

Total retained in pre-ART	**40,352**	**22%**
Started ART	**94,850**	**52%**
Defaulted	**44,778**	**25%**
Died	**2,011**	**1%**

Transfers between sites

Total not transferred out	**182,555**	**96%**
Transferred out	**6,742**	**4%**

* Subgroups may not add to 100% due to rounding

HTC site report

Malawi (national)

2015 Q1 (1st month of quarter, 2nd month of quarter, 3rd month of quarter)

Clients at health facility (static)

HTC client details

*

Total HTC clients served		
Total HIV tested	**471,270**	**100%**
Sex		
Males tested	**153,894**	**33%**
Females tested	**317,376**	**67%**
Females non-pregnant	**155,822**	49%
Females pregnant	**161,554**	51%
Age		
Children 0-14 yrs	**44,151**	**9%**
Children below 12 mths (Age group A)	**2,365**	5%
Children 12 mths - 14 yrs (Age group B)	**41,786**	95%
Adults 15+ years	**427,119**	**91%**
Young adults 15-24 years (Age group C)	**195,184**	46%
Older adults 25+ yrs (Age group D)	**231,935**	54%
HTC access type		
PITC	**275,409**	**58%**
Family Referral Slip (FRS)	**1,602**	**0%**
Other (VCT, etc.) HTC access	**194,259**	**41%**
HTC first time / repeat		
Never tested before	**147,145**	**31%**
Previously accessed HTC	**324,125**	**69%**
Last negative	**311,548**	96%
Last positive	**10,157**	3%
Last exposed infant	**2,067**	1%
Last inconclusive	**353**	0%
Counseling session type / Partner present		
Counseled with partner / partner present	**112,524**	**24%**
Counseled alone / Partner not present	**358,746**	**76%**
Outcome summary (HIV test)		
Single test negative	**429,863**	**91%**
Single test positive	**150**	**0%**
Test 1&2 negative	**489**	**0%**
Test 1&2 positive	**39,495**	**8%**
Test 1&2 discordant	**1,273**	**0%**
Final result given to client		
Results among clients never tested / last negative	**461,408**	**98%**
New negative	**429,364**	93%
New positive	**29,843**	6%
New exposed infants	**917**	0%
New inconclusive	**1,284**	0%
Confirmatory results (previous positive clients)	**9,862**	**2%**
Confirmatory positive	**9,295**	94%
Confirmatory inconclusive	**567**	6%

* Subgroups may not add to 100% due to rounding

HTC client details *

Partner / Family HTC referral slips

Sum of slips given		**13,437**	**100%**
	Total clients presenting with referral slip	**1,602**	12%
	Total failed referrals (slips not returned)	**11,835**	88%

Clients tested in the community

HTC client details *

Total HTC clients served

Total HIV tested		**12,796**	**100%**

Sex

Males tested		**5,047**	**39%**
Females tested		**7,749**	**61%**
	Females non-pregnant	**5,317**	69%
	Females pregnant	**2,432**	31%

Age

Children 0-14 yrs		**2,388**	**19%**
	Children below 12 mths (Age group A)	**10**	0%
	Children 12 mths - 14 yrs (Age group B)	**2,378**	100%
Adults 15+ years		**10,408**	**81%**
	Young adults 15-24 years (Age group C)	**6,346**	61%
	Older adults 25+ yrs (Age group D)	**4,062**	39%

HTC access type

PITC		**3,586**	**28%**
Family Referral Slip (FRS)		**3**	**0%**
Other (VCT, etc.) HTC access		**9,207**	**72%**

HTC first time / repeat

Never tested before		**5,535**	**43%**
Previously accessed HTC		**7,261**	**57%**
	Last negative	**7,197**	99%
	Last positive	**55**	1%
	Last exposed infant	**6**	0%
	Last inconclusive	**3**	0%

Counseling session type / Partner present

Counseled with partner / partner present		**1,424**	**11%**
Counseled alone / Partner not present		**11,372**	**89%**

Outcome summary (HIV test)

Single test negative		**12,468**	**97%**
Single test positive		**4**	**0%**
Test 1&2 negative		**4**	**0%**
Test 1&2 positive		**313**	**2%**
Test 1&2 discordant		**7**	**0%**

HTC site report

Malawi (national)

2015 Q1 (1st month of quarter, 2nd month of quarter, 3rd month of quarter)

HTC client details

*

Final result given to client

Results among clients never tested / last negative		**12,749**	**100%**
	New negative	**12,466**	98%
	New positive	**277**	2%
	New exposed infants	**3**	0%
	New inconclusive	**3**	0%
Confirmatory results (previous positive clients)		**47**	**0%**
	Confirmatory positive	**42**	89%
	Confirmatory inconclusive	**5**	11%

Partner / Family HTC referral slips

Sum of slips given		**136**	**100%**
	Total clients presenting with referral slip	**3**	2%
	Total failed referrals (slips not returned)	**133**	98%

Clients at stand-alone HTC sites

HTC client details

*

Total HTC clients served

Total HIV tested		**16,293**	**100%**

Sex

Males tested		**7,854**	**48%**
Females tested		**8,439**	**52%**
	Females non-pregnant	**7,016**	83%
	Females pregnant	**1,423**	17%

Age

Children 0-14 yrs		**373**	**2%**
	Children below 12 mths (Age group A)	**5**	1%
	Children 12 mths - 14 yrs (Age group B)	**368**	99%
Adults 15+ years		**15,920**	**98%**
	Young adults 15-24 years (Age group C)	**5,954**	37%
	Older adults 25+ yrs (Age group D)	**9,966**	63%

HTC access type

PITC		**8,401**	**52%**
Family Referral Slip (FRS)		**97**	**1%**
Other (VCT, etc.) HTC access		**7,795**	**48%**

HTC first time / repeat

Never tested before		**4,400**	**27%**
Previously accessed HTC		**11,893**	**73%**
	Last negative	**11,519**	97%
	Last positive	**362**	3%
	Last exposed infant	**8**	0%
	Last inconclusive	**4**	0%

Counseling session type / Partner present

Counseled with partner / partner present		**1,523**	**9%**
Counseled alone / Partner not present		**14,770**	**91%**

* Subgroups may not add to 100% due to rounding

HTC site report

Malawi (national)

2015 Q1 (1st month of quarter, 2nd month of quarter, 3rd month of quarter)

HTC client details

*

Outcome summary (HIV test)

Single test negative	**14,526**	**89%**
Single test positive	**1**	**0%**
Test 1&2 negative	**8**	**0%**
Test 1&2 positive	**1,730**	**11%**
Test 1&2 discordant	**28**	**0%**

Final result given to client

Results among clients never tested / last negative		**16,210**	**99%**
	New negative	**14,535**	90%
	New positive	**1,371**	8%
	New exposed infants	**6**	0%
	New inconclusive	**298**	2%
Confirmatory results (previous positive clients)		**83**	**1%**
	Confirmatory positive	**77**	93%
	Confirmatory inconclusive	**6**	7%

Partner / Family HTC referral slips

Sum of slips given		**812**	**100%**
	Total clients presenting with referral slip	**97**	12%
	Total failed referrals (slips not returned)	**715**	88%

* Subgroups may not add to 100% due to rounding

Maternity

Malawi (national)

2015 Q1 (1st month of quarter, 2nd month of quarter, 3rd month of quarter)

Maternal details

*

Admissions in the reporting period

Total admissions (referrals double-counted)	**129,368**	**100%**
Not referred to other site (total women)	**122,766**	95%
Referred out before delivery (multiple admissions)	**6,602**	5%

HIV status ascertainment

HIV status not ascertained	**4,987**	**4%**
HIV status ascertained	**120,651**	**96%**
Valid previous test result	**118,081**	98%
Previous negative	**109,161**	92%
Previous positive	**8,920**	8%
New test at maternity	**2,570**	2%
New negative	**2,348**	91%
New positive	**222**	9%

HIV status summary

Total women HIV negative	**111,509**	**92%**
Total women HIV positive	**9,142**	**8%**

ARVs during pregnancy (among HIV pos)

No ARV in pregnancy	**282**	**3%**
Any ARVs	**8,860**	**97%**
ART (by time of initiation)	**8,860**	100%
ART initiated before pregnancy	**6,596**	74%
ART initiated in 1st / 2nd trimester	**1,164**	13%
ART initiated in 3rd trimester	**902**	10%
ART initiated during labour	**198**	2%

Obstetric complications

No obstetric complications	**110,517**	**88%**
Any obstetric complications	**15,121**	**12%**
Haemorrhage	**2,475**	16%
Haemorrhage ante-partum	**805**	33%
Haemorrhage post-partum	**1,670**	67%
Obstr / prol labour	**4,840**	32%
(pre-) Eclampsia	**1,136**	8%
Maternal sepsis	**160**	1%
Ruptured uterus	**128**	1%
Other obstetric complications	**6,382**	42%

Emergency obstetric care

Oxytocin	**125,115**	**96%**
Anticonvulsive	**542**	**0%**
Antibiotics	**4,615**	**4%**
Blood transfusion	**298**	**0%**
Manual removal of placenta	**409**	**0%**

Vitamin A

Vit A not given	**39,020**	**31%**
Vit A given	**86,618**	**69%**

* Subgroups may not add to 100% due to rounding

Maternity

Malawi (national)

2015 Q1 (1st month of quarter, 2nd month of quarter, 3rd month of quarter)

Maternal details

*

Staff conducting delivery

Category A: MO, CO, nurse/midwife, MA	**112,425**	**94%**
Category B: PA, WA, HSA	**585**	**0%**
Category C: Other	**6,026**	**5%**

Mother survival

Mother alive	**118,944**	**100%**
Mother died	**92**	**0%**

Infant details

*

Single babies / multiple deliveries

Total babies delivered	**121,153**	**100%**
Single babies	**117,192**	97%
Twin / multiple babies	**3,961**	3%

Delivery place

Total deliveries at a health facility	**114,970**	**95%**
This facility	**114,594**	100%
Other facility	**376**	0%
Total deliveries before reaching the facility	**6,183**	**5%**
In transit	**4,025**	65%
Home / TBA	**2,158**	35%

Delivery mode

Spontaneous vaginal	**109,243**	**90%**
Vacuum extraction	**1,473**	**1%**
Breech	**2,189**	**2%**
Caesarean section	**8,248**	**7%**

Infant complications

No infant complications	**104,794**	**86%**
Total infants with complications	**16,359**	**14%**
Prematurity	**3,773**	23%
Weight less 2500g	**4,961**	30%
Asphyxia	**4,790**	29%
Sepsis	**1,001**	6%
Other newborn complication	**1,834**	11%

Infant survival

Total live births	**119,103**	**98%**
Discharged alive	**118,003**	99%
Neonatal deaths	**1,100**	1%
Stillbirths	**2,050**	**2%**
Stillbirth, fresh	**1,112**	54%
Stillbirth, macerated	**938**	46%

Maternity

Malawi (national)

2015 Q1 (1st month of quarter, 2nd month of quarter, 3rd month of quarter)

Infant details

*

HIV exposure / ARV proph. (among discharged alive)

Infants with unknown HIV exposure status	**4,112**	**3%**
Infants with known HIV exposure status	**113,891**	**97%**
Not HIV exposed	**105,392**	93%
HIV exposed	**8,499**	7%
Received no ARVs	**633**	7%
Received ARVs	**7,866**	93%
Nevirapine	**7,866**	100%

Breastfeeding initiated

BF not started within 60min	**10,177**	**8%**
BF started within 60min	**110,976**	**92%**

Tetracycline eye ointment given

TO not given	**20,710**	**17%**
TO given	**100,443**	**83%**

* Subgroups may not add to 100% due to rounding

STI site report

Malawi (national)

2015 Q1 (1st month of quarter, 2nd month of quarter, 3rd month of quarter)

STI clients treated in the reporting period

*

Total STI clients

Total STI clients treated		**54,897**	**100%**
	Index patients treated (symptomatic)	**44,127**	80%
	Partners treated	**10,770**	20%

Sex

Males		**21,850**	**40%**
Females		**33,047**	**60%**
	Non-pregnant	**28,809**	87%
	Pregnant	**4,238**	13%

Age group

Age group A (0-19 years)		**4,898**	**9%**
Age group B (20-24 years)		**13,176**	**24%**
Age group C (25+ years)		**36,823**	**67%**

Client type

Symptomatic cases		**48,409**	**88%**
	Index cases	**44,127**	91%
	Partners symptomatic	**4,282**	9%
Partners asymptomatic		**6,488**	**12%**

STI treatment history

Never treated for STI		**40,691**	**74%**
Previously treated for STI		**14,206**	**26%**
	Old >3 months ago	**9,608**	68%
	Recent ≤3 months ago	**4,598**	32%

STI syndromic diagnosis

GUD		**10,229**	**17%**
UD		**13,008**	**22%**
AVD		**17,974**	**30%**
	Low risk	**7,219**	40%
	High risk	**10,755**	60%
LAP		**9,246**	**15%**
SS		**753**	**1%**
BU		**694**	**1%**
BA		**1,310**	**2%**
NC		**274**	**0%**
Genital Warts		**612**	**1%**
Syphilis RPR VDRL		**1,603**	**3%**
Other STI		**4,215**	**7%**

STI partner notification

Total partner notification slips issued		**19,738**	**100%**
	Total partners returned	**10,770**	55%
	Total partners not seen	**8,968**	45%

* Subgroups may not add to 100% due to rounding

ART survival analysis

Malawi (national)

2015 Q1 (Quarter)

12 month survival children

Survival and retention in ART program *

ART cohort registration group outcomes

Total ART clinic registrations	**2,651**	**100%**
Transfers out (double counted)	**249**	9%
Total not transferred out (patients in cohort)	**2,402**	91%
Total alive on ART	**1,859**	77%
Total not retained	**543**	23%
Defaulted	**423**	78%
Stopped ART	**14**	3%
Died	**106**	20%

12 month survival all ages

Survival and retention in ART program *

ART cohort registration group outcomes

Total ART clinic registrations	**31,492**	**100%**
Transfers out (double counted)	**2,834**	9%
Total not transferred out (patients in cohort)	**28,658**	91%
Total alive on ART	**21,827**	76%
Total not retained	**6,831**	24%
Defaulted	**5,712**	84%
Stopped ART	**54**	1%
Died	**1,065**	16%

24 month survival all ages

Survival and retention in ART program *

ART cohort registration group outcomes

Total ART clinic registrations	**33,735**	**100%**
Transfers out (double counted)	**4,357**	13%
Total not transferred out (patients in cohort)	**29,378**	87%
Total alive on ART	**21,136**	72%
Total not retained	**8,242**	28%
Defaulted	**6,595**	80%
Stopped ART	**108**	1%
Died	**1,539**	19%

36 month survival all ages

Survival and retention in ART program *

ART cohort registration group outcomes

Total ART clinic registrations	**43,137**	**100%**
Transfers out (double counted)	**6,962**	16%
Total not transferred out (patients in cohort)	**36,175**	84%
Total alive on ART	**25,610**	71%
Total not retained	**10,565**	29%
Defaulted	**8,146**	77%
Stopped ART	**131**	1%
Died	**2,288**	22%

* Subgroups may not add to 100% due to rounding

ART survival analysis

Malawi (national)

2015 Q1 (Quarter)

48 month survival all ages

Survival and retention in ART program *

ART cohort registration group outcomes

Total ART clinic registrations	**22,731**	**100%**
Transfers out (double counted)	**5,512**	24%
Total not transferred out (patients in cohort)	**17,219**	76%
Total alive on ART	**11,221**	65%
Total not retained	**5,998**	35%
Defaulted	**4,019**	67%
Stopped ART	**74**	1%
Died	**1,905**	32%

60 month survival all ages

Survival and retention in ART program *

ART cohort registration group outcomes

Total ART clinic registrations	**23,410**	**100%**
Transfers out (double counted)	**5,950**	25%
Total not transferred out (patients in cohort)	**17,460**	75%
Total alive on ART	**10,932**	63%
Total not retained	**6,528**	37%
Defaulted	**4,423**	68%
Stopped ART	**87**	1%
Died	**2,018**	31%

72 month survival all ages

Survival and retention in ART program *

ART cohort registration group outcomes

Total ART clinic registrations	**21,100**	**100%**
Transfers out (double counted)	**5,596**	27%
Total not transferred out (patients in cohort)	**15,504**	73%
Total alive on ART	**9,166**	59%
Total not retained	**6,338**	41%
Defaulted	**4,007**	63%
Stopped ART	**54**	1%
Died	**2,277**	36%

84 month survival all ages

Survival and retention in ART program *

ART cohort registration group outcomes

Total ART clinic registrations	**16,923**	**100%**
Transfers out (double counted)	**4,797**	28%
Total not transferred out (patients in cohort)	**12,126**	72%
Total alive on ART	**6,656**	55%
Total not retained	**5,470**	45%
Defaulted	**3,230**	59%
Stopped ART	**63**	1%
Died	**2,177**	40%

* Subgroups may not add to 100% due to rounding

96 month survival all ages

Survival and retention in ART program *

ART cohort registration group outcomes

Total ART clinic registrations	**12,909**	**100%**
Transfers out (double counted)	**3,645**	28%
Total not transferred out (patients in cohort)	**9,264**	72%
Total alive on ART	**4,624**	50%
Total not retained	**4,640**	50%
Defaulted	**2,441**	53%
Stopped ART	**73**	2%
Died	**2,126**	46%

108 month survival all ages

Survival and retention in ART program *

ART cohort registration group outcomes

Total ART clinic registrations	**9,475**	**100%**
Transfers out (double counted)	**2,808**	30%
Total not transferred out (patients in cohort)	**6,667**	70%
Total alive on ART	**3,073**	46%
Total not retained	**3,594**	54%
Defaulted	**1,713**	48%
Stopped ART	**36**	1%
Died	**1,845**	51%

120 month survival all ages

Survival and retention in ART program *

ART cohort registration group outcomes

Total ART clinic registrations	**4,512**	**100%**
Transfers out (double counted)	**1,294**	29%
Total not transferred out (patients in cohort)	**3,218**	71%
Total alive on ART	**1,479**	46%
Total not retained	**1,739**	54%
Defaulted	**867**	50%
Stopped ART	**38**	2%
Died	**834**	48%

6 month survival OptionB+

Survival and retention in ART program *

ART cohort registration group outcomes

Total ART clinic registrations	**8,508**	**100%**
Transfers out (double counted)	**472**	6%
Total not transferred out (patients in cohort)	**8,036**	94%
Total alive on ART	**6,148**	77%
Total not retained	**1,888**	23%
Defaulted	**1,835**	97%
Stopped ART	**12**	1%
Died	**41**	2%

ART survival analysis

Malawi (national)

2015 Q1 (Quarter)

12 month survival OptionB+

Survival and retention in ART program *

ART cohort registration group outcomes

Total ART clinic registrations		**8,392**	**100%**
	Transfers out (double counted)	**664**	8%
	Total not transferred out (patients in cohort)	**7,728**	92%
	Total alive on ART	**5,601**	72%
	Total not retained	**2,127**	28%
	Defaulted	**2,029**	95%
	Stopped ART	**32**	2%
	Died	**66**	3%

24 month survival OptionB+

Survival and retention in ART program *

ART cohort registration group outcomes

Total ART clinic registrations		**8,714**	**100%**
	Transfers out (double counted)	**863**	10%
	Total not transferred out (patients in cohort)	**7,851**	90%
	Total alive on ART	**5,411**	69%
	Total not retained	**2,440**	31%
	Defaulted	**2,314**	95%
	Stopped ART	**42**	2%
	Died	**84**	3%

36 month survival OptionB+

Survival and retention in ART program *

ART cohort registration group outcomes

Total ART clinic registrations		**11,951**	**100%**
	Transfers out (double counted)	**1,466**	12%
	Total not transferred out (patients in cohort)	**10,485**	88%
	Total alive on ART	**7,315**	70%
	Total not retained	**3,170**	30%
	Defaulted	**2,916**	92%
	Stopped ART	**53**	2%
	Died	**201**	6%

* Subgroups may not add to 100% due to rounding

STI site report

Malawi (national)

2015 Q1 (1st month of quarter, 2nd month of quarter, 3rd month of quarter)

STI clients treated in the reporting period

*

HIV test / ART status

HIV status not ascertained	**28,580**	**52%**
HIV status ascertained	**26,317**	**48%**
HIV negative (new test)	**18,730**	71%
HIV positive	**7,587**	29%
New positive	**1,607**	21%
Previous positive	**5,980**	79%
Not on ART	**1,358**	23%
On ART	**4,622**	77%

STI clients referred for services

Lab	**567**	**3%**
Gynae review	**387**	**2%**
Surgical review	**561**	**3%**
Repeat HTC	**17,766**	**83%**
ART (for assessment)	**856**	**4%**
PMTCT	**130**	**1%**
Other (service referrals)	**1,025**	**5%**

* Subgroups may not add to 100% due to rounding